



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Goat anti Rat alpha-2 macroglobulin

Catalogue number: **GARa/A2M**

Clone	Polyclonal
Product Type	Primary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Rat
Application	Immunoprecipitation

### Background

The reactivity of the antiserum is restricted to rat alpha-2 macroglobulin. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal rat plasma a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified alpha-2 macroglobulin. No reaction is obtained with any other plasma protein components or serum. In precipitating techniques as immunoelectrophoresis and single or double radial immunodiffusion (Mancini, Ouchterlony) to identify the presence of alpha-2 macroglobulin in rat plasma or other body fluids or to determine its concentration. This antiserum is not intended for use in non-precipitating antibody binding or other highly sensitive assays, but this does exclude such use if proper controls are included.

### Source

Alpha-2 macroglobulin (clotting factor I) is the major protein in the alpha-2 region of rat serum after separation by immunoelectrophoresis. The concentration in normal plasma is about 2.5 mg/ml. It is a large polymer and a strong antigen. It forms about one third in the total of the alpha-2 globulins. The protein is isolated by ammonium sulphate precipitation, ion exchange chromatography, gel filtration and preparative starch block electrophoresis. In the purification the activation of kallikrein is avoided, since it forms an irreversible bond with alpha-2 macroglobulin, thus contaminating the final immunogen. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

Delipidated, heat inactivated, lyophilized, stable whole antiserum  
No preservative added as it may interfere with the antibody

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

activity. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal goat serum. No foreign proteins added.

#### **Applications**

Immunoprecipitation. The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.

#### **Cross Reactivity**

The antiSerum does not cross-react with any other component of Rat plasma. Inter-species cross-reactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail, however in double radial immunodiffusion a reaction has been observed with Mouse.

#### **Specificity**

Precipitating polyclonal Goat antiSerum to Rat alpha-2 macroglobulin.

#### **Storage**

Lyophilized at +4° C--at least 10 years. Reconstituted at or below -20° C--3-5 years. Reconstituted at +4° C--7 days

#### **Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.