



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Goat anti Rat C3c, conjugated with FITC

Catalogue number: **GARa/C3c/FITC**

Clone	Polyclonal
Product Type	Primary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Rat
Application	ELISA Immunocytochemistry Immunofluorescence Immunohistochemistry (frozen)

Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

Background

In immunoelectrophoresis against fresh rat serum, a single precipitin line is obtained in the beta-1 region representing native C3. Against serum containing partly activated C3, a precipitin line is obtained which extends from the beta-1 into the alpha-2 region, demonstrating a gradient. In old serum containing totally activated C3 a single precipitin line in the alpha-2 region is obtained. Antisera to C3c can also react with the fragments C3b, C3bi and smaller fragments, since they all carry antigenic determinants of the C3c domain. The product does not react with any other proteins component of rat serum or plasma. The fluorescent immunoconjugate to rat C3c is used to determine the presence and pattern of C3 in tissue lesions using immunohistochemical staining techniques. Locally deposited immune complexes in tissue usually contain complement, pointing to activation of the classical pathway. Complement activation in vivo implies active disease and may contribute to the elicitation of the pathogenesis and the extent of tissue destruction. Sometimes the diagnosis can be based directly on laboratory findings. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Source

C3 is the most abundant complement protein in rat serum. Its biological function strongly resembles that of C3 in man and other laboratory animal species. It has a central role in the activation system being common to both pathways. Activation of C3 is

achieved by very specific limited proteolysis resulting in the release of a number of degradation fragments. The anaphylotoxin C3a promotes smooth muscle contraction and increases vascular permeability; the large C3b fragment is involved in binding to the complement activator and can interact with specific receptors to allow efficient clearance of the activating cell or particle; degradation fragments of C3b (C3bi, C3c, C3dg, C3d) are important in receptor binding and clearance mechanisms, in virus neutralization and possibly in the immune response. The antiserum is raised against C3c, which is the major fragment resulting from C3 cleavage by C3 convertase and factor I. It is composed of an intact beta chain bound to two fragments of the alpha chain. Consequently the antiserum reacts with both native and activated C3. It may also react with the fragments C3b, C3bi and C3dg, since they all carry antigenic epitopes of the C3c domain. C3c is isolated and purified from pooled normal rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure

Product

Fluorochrome-coupled purified hyperimmune IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2) No preservative added, as it may interfere with the antibody activity.

Applications

ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.

Cross Reactivity

The antiSerum does not cross-react with any other component of Rat plasma. Inter-species cross-reactivity is a normal feature of antibodies to plasma proteins since they frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.

Specificity

Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal Goat antiSerum to C3c fragment of Rat complement factor C3.

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as

reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.

[Home](#) | [Company Profile](#) | [Catalogue](#) | [Distributors](#) | [Contact](#) |

Content: Nordic-MUbio BV - Copyright © 2015