



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Goat anti Rat IgG1 (subclass specific), conjugated with FITC

Catalogue number: **GARa/IgG1/FITC**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Goat
Species reactivity	Rat
Application	ELISA Immunocytochemistry Immunofluorescence Immunohistochemistry (frozen)

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

### Background

reactivity of the antiserum is directed to the subclass IgG1 as tested against rat sera and a panel of purified homogenous immunoglobulins. It does not react with other subclasses of IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein in rat serum, as tested by immunoelectrophoresis and double radial immunodiffusion. Further tested on the level of sensitivity of direct and indirect immunofluorescence staining methods applied to lymphoid cells. When used on vital peripheral blood cells this immunoconjugate does not react with Fc receptors present on cell surface membranes. To identify and measure IgG1, antigen or antibody, at the cellular and subcellular level by immunofluorescence staining of appropriately treated cell and tissue substrates, and to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of rat origin in the middle layer of the indirect test procedure. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

### Source

Pools of purified homogenous IgG1 isolated from pooled rat serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

Fluorochrome-coupled purified hyperimmune goat IgG lyophilized

from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

#### **Applications**

ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.

#### **Cross Reactivity**

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this product has not been tested in detail, however in double radial immunodiffusion a weak reaction with Mouse has been observed.

#### **Specificity**

Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal Goat antiSerum to Rat IgG1, subclass specific.

#### **Storage**

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate. Lyophilized at +4° C--at least 10 years. Reconstituted at or below -20° C--3-5 years. Reconstituted at +4° C--7 days

#### **Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.