



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Goat IgG, conjugated with FITC

Catalogue number: **GlgG/FITC**

Product Type	Isotype control
Units	1 ml
Host	Goat
Species reactivity	Goat
Application	ELISA Immunocytochemistry Immunofluorescence Immunohistochemistry (frozen)

Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

Background

As an efficient general counterstain for cell and tissue substrates with immunoconjugates with a label contrasting with the greenish-yellow fluorescence of this product. The purpose of counterstaining is to enhance the optical contrast of the immunospecific staining against a counterstained background.

Product

Fluorochrome coupled purified goat IgG in phosphate buffered saline (pH7.2). No preservative added

Applications

ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence. The conjugate is shipped at ambient temperature and may be stored at +4°C, prolonged storage at -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Spin down to remove insoluble particles, divide into small aliquots, and store frozen at or below -20°C. Prior to use an aliquot is thawed at ambient temperature, spun down again, and used to prepare working dilutions by adding sterile PBS (pH7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the conjugate. Diluted product should be stored at +4°C, not refrozen, and preferably used the same day.

Specificity

Fluorescein isothiocyanate-conjugated polyclonal Goat IgG

Storage

Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years reconstituted at +4°C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.