



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



---

## Mouse anti Myeloperoxidase-C2 (MPO-C2), conjugated to PE

---

[nordicmubio.com/products/Mouse-anti-Myeloperoxidase-C2-MPO-C2-conjugated-to-PE/GM-4193](https://nordicmubio.com/products/Mouse-anti-Myeloperoxidase-C2-MPO-C2-conjugated-to-PE/GM-4193)

Catalog number: **GM-4193**

Clone	8E6
Isotype	IgG1
Product Type	Primary Antibodies
Units	2ml (100 Tests)
Host	Mouse
Species Reactivity	Human
Application	Flow Cytometry

### Background

Myeloperoxidase (MPO) is a glycoprotein present in the azurophil (primary) granules of myeloid cells, which appears in the myeloblast stage of myeloid cell differentiation. MPO is the most common functional protein of myeloid cells and is involved in the inflammatory response. It helps to kill microbes by breaking down peroxide in the presence of halide ions, contributing to the bactericidal function of granulocytes. The primary translation product of MPO undergoes glycosylation with production of the 89 kDa heme-free apopro-MPO form followed by incorporation of heme and conversion into the enzymatically active pro-MPO form. Subsequently, pro-MPO becomes targeted to azurophil granules where final processing occurs to produce mature dimeric MPO consisting of the 59-64 kDa MPO  $\alpha$ -chain and the 14 kDa MPO chain. The MPO-C2 mAb (clone 8E6) recognizes virtually all myelomonocytic cells including AML blasts. The MPO-C2 mAb permits the identification and enumeration of myeloid cells in normal and malignant human blood and bone marrow using flow cytometry. Results must be interpreted by a certified professional before final interpretation. Analyses performed with this antibody should be paralleled by positive and negative controls. If unexpected results are obtained which cannot be attributed to differences in laboratory procedures, please contact us.

## **Product**

2 ml of PE-conjugated anti Myeloperoxidase-C2 (clone 8E6) in PBS pH 7.2, 1% BSA, and 0.05% NaN<sub>3</sub>, approximately 100 tests.

*Product Form:* PE

*Formulation:* PBS pH 7.2, 1% BSA, 0.05% NaN<sub>3</sub>

*Purification Method:* Purified by Chromatography

## **Specificity**

The MPO-C2 mAb (clone 8E2) reacts with human myeloperoxidase (MPO) expressed by normal and malignant myelomonocytic cells. The sensitivity of MPO-C2 mAb is determined by staining well-defined blood samples from representative donors with serial-fold mAb dilutions to obtain a titration curve that allows relating the mAb concentration to the percentage of stained cells and geometric MFI (mean fluorescence intensity). For this purpose, a mAb-concentration range is selected to include both the saturation point (i.e. the mAb dilution expected to bind all epitopes on the target cell) and the detection threshold (i.e. the mAb dilution expected to represent the least amount of mAb needed to detect an identical percentage of cells). In practice, 50 µl of leukocytes containing 10<sup>7</sup> cells/ml are stained with 20 µl mAb of various dilutions to obtain a titration curve and to identify the saturation point and detection threshold. The final concentration of the product is then adjusted to be at least 3-fold above the detection threshold. In addition and to control lot-to-lot variation, the given lot is compared and adjusted to fluorescence standards with defined intensity.

## **Applications**

Permeabilization and Staining Procedure - In combination with our Permeabilization Kit FIX&PERM® (Cat. No. GAS-002) intracellular MPO can be easily stained in cell suspensions. - For each sample to be analyzed add 50 µl of whole blood, bone marrow or mononuclear cell suspension in a 5 ml tube - Add 100 µl of Reagent A (Fixation Medium, stored and used at room temperature) - Incubate for 15 minutes at room temperature - Add 5 ml phosphate buffered saline and centrifuge cells for 5 minutes at 300 g - Remove supernatant and add to cell pellet 100 µl Reagent B (Permeabilization Medium) and 20 µl of the MPO-C2 monoclonal antibody conjugate - Vortex at low speed for 1-2 seconds - Incubate for 15 minutes at room temperature - Wash cells with phosphate buffered saline as described above - Remove supernatant and resuspend cells in sheath fluid for immediate analysis or resuspend cells in 0.5 ml 1.0 % formaldehyde and store them at 2-8°C in the dark. Analyze fixed cells within 24 hours.

## **Storage**

Nordic-MUbio monoclonal antibody reagents contain optimal concentrations of affinity-purified antibody. For stability reasons this monoclonal antibody solution contains sodium azide. These reagents should be stored at 2-8°C (DO NOT FREEZE!) and protected from prolonged exposure to light. If a slight precipitation occurs upon storage, this should be

removed by centrifugation. It will not affect the performance or the concentration of the product. Stability of the reagent: Please refer to the expiry date printed onto the vial. The use of the reagent after the expiration date is not recommended.

*Shipping Conditions:* Ship at ambient temperature

### **Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. It may contain hazardous ingredients. Please refer to the Safety Data Sheets (SDS) for additional information and proper handling procedures. Dispose product remainders according to local regulations. This datasheet is as accurate as reasonably achievable, but Exalpa Biologicals accepts no liability for any inaccuracies or omissions in this information.

### **References**

1. Andersson, E., Hellman, L., Gullberg, U. & Olsson, I. (1998) *J Biol Chem* 273, 4747-53.
2. Catovsky, D., Matutes, E., Buccheri, V., Shetty, V., Hanslip, J., Yoshida, N. & Morilla, R. (1991) *Ann Hematol* 62, 16-21.
3. Cramer, E., Pryzwansky, K. B., Villeval, J. L., Testa, U. & Breton-Gorius, J. (1985) *Blood* 65, 423-32.
4. Gullberg, U., Andersson, E., Garwicz, D., Lindmark, A. & Olsson, I. (1997) *Eur J Haematol* 58, 137-53.
5. Imamura, N. (1998) *Am J Hematol* 58, 241-3.
6. Knapp, W., Majdic, O. & Strobl, H. (1993) *Recent Results Cancer Res* 131, 31-40.
7. Koeffler, H. P., Ranyard, J. & Pertcheck, M. (1985) *Blood* 65, 484-91.
8. Lanza, F., Latorraca, A., Moretti, S., Castagnari, B., Ferrari, L. & Castoldi, G. (1997) *Cytometry* 30, 134-44.
9. Murao, S., Stevens, F. J., Ito, A. & Huberman, E. (1988) *Proc Natl Acad Sci U S A* 85, 1232-6.
10. Nakase, K., Sartor, M. & Bradstock (1998) *Cytometry* 34, 198-202.
11. Nauseef, W. M. (1990) *Hematol Pathol* 4, 165-78.
12. Nauseef, W. M., Olsson, I. & Arnljots, K. (1988) *Eur J Haematol* 40, 97-110.
13. Srivastava, C. H., Rado, T. A., Bauerle, D. & Broxmeyer, H. E. (1991) *J Immunol* 146, 1014-9.
14. Strobl, H. & Knapp, W. (2004) *J Biol Regul Homeost Agents* 18, 335-9.
15. Strobl, H., Takimoto, M., Majdic, O., Fritsch, G., Scheinecker, C., Hocker, P. & Knapp, W. (1993) *Blood* 82, 2069-78.
16. Tsuruta, T., Tani, K., Hoshika, A. & Asano, S. (1999) *Leuk Lymphoma* 32, 257-67.
17. van der Schoot, C. E., Daams, G. M., Pinkster, J., Vet, R. & von dem Borne, A. E. (1990) *Br J Haematol* 74, 173-8.
18. van der Schoot, C. E., von dem Borne, A. E. & Tetteroo, P. A. (1987) *Acta Haematol* 78 Suppl 1, 32-40.
19. Zaki, S. R., Austin, G. E., Swan, D., Srinivasan, A., Ragab, A. H. & Chan, W. C. (1989) *Blood* 74, 2096-102.

### **Warranty**

The products sold hereunder are warranted only to conform to the quantity and contents stated on the label at the time of delivery to the customer. There are no warranties, expressed or implied, that extend beyond the description on the label of the product. Exalpa's sole liability is limited to either replacement of the products or refund of the purchase price. Exalpa is not liable for property damage, personal injury, or economic loss caused by the product.

**Safety Datasheet(s) for this product:**

NM\_Sodium Azide