

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



Mouse anti Human Bence Jones kappa (surface and hidden determinants), conjugated with Biotin

Catalogue number: MAHu/BJK/Bio

Clone	NI 285, NI 250
lsotype	lgG1
Product Type	Secondary Antibodies
Units	0.2 mg
Host	Mouse
Species reactivity	Human
Application	Dot blot ELISA Immunoblotting Immunocytochemistry Immunohistochemistry (paraffin)

Background

The reactivity of this preparation of two monoclonal antibodies is restricted to a surface determinant on the kappa light chain. It is reacting with polyclonal and monoclonal immunoglobulins of the kappa type, as well as free kappa light chains as tested in direct binding enzyme immunoassay, immunoblotting, immunoprecipitation and direct immunoperoxidase staining. To identify the light chain type of immunoglobulins or free light chains in human serum, other body fluids, cell and tissue substrates and to determine its concentration in techniques as ELISA, direct immunoperoxidase staining and immunoblotting. The optimum working dilution is an assay-related characteristic and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:100 to 1:500; in ELISA from 1:1,000 upwards; in Western blotting from 1:2.000 upwards. These data should be interpreted as general recommendations only.

Source

Highly purified Bence Jones kappa proteins isolated from pooled human urine.

Product

No preservative added, as it may interfere with the antibody activity. No foreign protein added.

Distributors

For Purchasing Information, please contact your local distributor

Find Distributor

Applications

ELISA,Immunocytochemistry,Immunohistochemistry (paraffin),Dot blot,Immunoblotting.

Cross Reactivity

The antibody does not react with any other component of the Human immunoglobulin system or any other Human plasma protein as tested. This antiSerum has not been tested for crossreactivity with other species.

Specificity

Biotin-conjugated purified monoclonal Mouse antibody to Human Ig kappa chain, free and bound

Storage

The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized product by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted ascites should be stored at +4°C, not refrozen, and preferably used the same day. Lyophilized at +4° C--at least 10 years. Reconstituted at or below -20° C--3-5 years. Reconstituted at +4° C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.

Home | Company Profile | Catalogue | Distributors | Contact

Content: Nordic-MUbio BV - Copyright © 2015