



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Mouse anti Human IgA2 (subclass specific), conjugated with FITC

Catalogue number: **MAHu/IgA2/FITC**

Clone	NI 512 (A89-038)
Isotype	IgG1
Product Type	Secondary Antibodies
Units	0.2 mg
Host	Mouse
Species reactivity	Human
Application	ELISA Immunocytochemistry Immunofluorescence Immunohistochemistry (frozen)

Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

Background

The reactivity of the antiserum is restricted to an subclass specific determinant on IgA2 molecule as tested in haemagglutination, haemagglutination inhibition, direct binding enzyme immunoassay, competitive inhibition enzyme immunoassay, immunoblotting, immunoprecipitation, latex agglutination assay and histochemistry (Results of an IUIS/WHO collaborative study, Mestecky J. et al. (1996) J. Immunol. Methods 193, 103-148). To identify the presence of IgA2 in human serum, other body fluids, cell and tissue substrates and to determine its concentration in techniques as immunofluorescence staining of cytoplasmic IgA2, and immunoblotting using a peroxidase labelled monoclonal antibody against FITC. The optimum working dilution is an assay-related characteristic. It may vary widely and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:20 to 1:100; in Western blotting from 1:200 upwards.

Source

Highly purified monoclonal IgA2 isolated from human serum.

Product

Purified monoclonal mouse IgG1 kappa conjugated with FITC, lyophilized from a solution in phosphate buffered saline (pH7.2). No preservative added, as it may interfere with the antibody activity. No foreign protein added.

Applications

ELISA, Immunocytochemistry, Immunohistochemistry (frozen),

(In)direct immunofluorescence.

Cross Reactivity

The antiSerum does not react with any other component of the Human Ig system or any other Human plasma protein as tested. This antiSerum has not been tested for cross-reactivity with other species.

Specificity

Fluorescein isothiocyanate-conjugated purified monoclonal Mouse antibody to Human IgA2, subclass specific

Storage

The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized product by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted solutions should be stored at +4°C, not refrozen, and preferably used the same day. Lyophilized at +4°C -- at least 10 years. Reconstituted at or below -20° C--3-5 years. Reconstituted at +4° C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.