

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Mouse anti Human secretory component (free and bound), conjugated with FITC

Catalogue number: MAHu/SC/FITC

Clone	NI 194-4 (A89-039)
Isotype	IgG1
Product Type	Secondary Antibodies
Units	0.2 mg
Host	Mouse
Species reactivity	Human
Application	ELISA
	Immunocytochemistry
	Immunofluorescence
	Immunohistochemistry (frozen)
1	

Distributors

For Purchasing Information, please contact your local distributor

Find Distributor

Background

The reactivity of the antiserum is restricted to determinants on the secretory component as tested in haemagglutination, haemagglutination inhibition, indirect binding enzyme immunoassay, competitive inhibition enzyme immunoassay, immunoblotting, immunoprecipitation, latex agglutination assay and histochemistry (Results of an IUIS/WHO collaborative study, Mestecky J. et al. (1996) J. Immunol. Methods 193, 103-148). To identify the presence of secretory component, free or bound in human milk, other body fluids, cell and tissue substrates and to determine its concentration in immunofluorescence staining techniques. The optimum working dilution is an assay-related characteristic and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:10 upwards. These data should be interpreted as general recommendations only.

Source

Highly purified secretory component isolated from human milk.

Product

Purified monoclonal mouse IgG1 kappa conjugated with fluorescein isothiocyanate, lyophilized from a solution in phosphate buffered saline (pH7.2). No preservative added, as it may interfere with the antibody activity. No foreign protein added.

Applications

ELISA,Immunocytochemistry,Immunohistochemistry (frozen), (In)direct immunofluorescence.

Cross Reactivity

The antiSerum does not react with any other component of the Human Ig system or any other plasma protein as tested. This antiSerum has not been tested for cross-reactivity with other species.

Specificity

Fluorescein isothiocyanate-conjugated purified monoclonal Mouse antibody to Human secretory component, free and bound.

Storage

The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized product by adding 0.5 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted solutions should be stored at +4°C, not refrozen, and preferably used the same day. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.

Home | Company Profile | Catalogue | Distributors | Contact

Content: Nordic-MUbio BV - Copyright © 2015