



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Mouse anti Human IgA1 (subclass specific)

nordicmubio.com/product/mouse-anti-human-iga1-subclass-specific-2

Catalogue number: **MAHu/IgA1/Mab**

| | |
|--------------|----------------------|
| Clone | NI 69-11 (A89-036) |
| Isotype | IgG1 |
| Product Type | Secondary Antibodies |
| Units | 0.2 mg |
| Host | Mouse |

Background

The reactivity of the antiserum is restricted to a subclass specific determinant on the CH2 domain of IgA1 as tested in haemagglutination, haemagglutination inhibition, direct binding enzyme immunoassay, competitive inhibition enzyme immunoassay, immunoblotting, immunoprecipitation, latex agglutination assay and histochemistry (Results of an IUIS/WHO collaborative study, Mestecky J. et al. (1996) J. Immunol. Methods 193, 103-148). To identify the presence of IgA1 in human serum, other body fluids, cell and tissue substrates and to determine its concentration in techniques as radio immuno assay, ELISA, indirect immunoperoxidase and immunofluorescence staining of cytoplasmic IgA1, haemagglutination and immunoblotting. The optimum working dilution is an assay-related characteristic and should always be determined by titration. For histochemical use optimum dilutions are mostly from 1:50 to 1:200; in ELISA from 1:200 upwards; in Western blotting from 1:500 upwards. These data should be interpreted as general recommendations only.

Source

Highly purified monoclonal IgA1 isolated from human serum.

Product

Purified mouse IgG1 kappa lyophilized from a solution in phosphate buffered saline (pH 7.2) IgG concentration is 0.4mg/ml. No foreign proteins added. Reconstitute the lyophilized ascites by adding 0.5 ml sterile distilled water.

Specificity

Purified monoclonal Mouse antibody to Human IgA1, subclass specific

Species Reactivity: The antiSerum does not react with any other component of the Human Ig system or any other Human plasma protein as tested. This antiSerum has not been tested for cross-reactivity with other species.

Applications

Indirect immunofluorescence,ELISA.

Storage

The lyophilized product is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Avoid repeated thawing and freezing. If a slight precipitation occurs upon storage, this should be removed by centrifugation and will not affect the performance of the product. Diluted solutions should be stored at +4°C, not refrozen, and preferably used the same day.

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.