



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Mouse anti Cytokeratin 10 / Keratin K10

 nordicmubio.com/products/mouse-anti-cytokeratin-10-keratin-k10/MUB0319P

Catalog number: **MUB0319P**

Clone	RKSE60
Isotype	IgG1
Product Type	Primary Antibodies
Units	0.1 mg
Host	Mouse
Species Reactivity	Canine Human Mouse Rat Swine Zebrafish
Application	Flow Cytometry Immunocytochemistry Immunohistochemistry (frozen) Western Blotting

Background

Cytokeratins are a subfamily of intermediate filament proteins and are characterized by a remarkable biochemical diversity, represented in Human epithelial tissues by at least 20 different polypeptides. They range in molecular weight between 40 kDa and 68 kDa and isoelectric pH between 4.9 – 7.8. The individual Human Cytokeratins are numbered 1 to 20. The various epithelia in the Human body usually express Cytokeratins which are not only characteristic of the type of epithelium, but also related to the degree of maturation or differentiation within an epithelium. Cytokeratin subtype expression patterns are used to an increasing extent in the distinction of different types of epithelial malignancies. The Cytokeratin antibodies are not only of assistance in the differential diagnosis of tumors using immunohistochemistry on tissue sections, but are also a useful tool in cytopathology and flow cytometric assays.

Source

RKSE60 is a Mouse monoclonal IgG1 antibody derived by fusion of SP2/0 Mouse myeloma cells with spleen cells from a BALB/c Mouse immunized with Cytokeratins from the Human epidermis.

Product

Each vial contains 100 ul 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.

Formulation: Each vial contains 100 ul 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.

Specificity

RKSE60 reacts exclusively with Cytokeratin 10 which is present in Keratinizing stratified epithelia and in differentiated areas of highly differentiated squamous cell carcinomas.

Applications

RKSE60 is suitable for immunoblotting, immunocytochemistry, immunohistochemistry on frozen tissues and flow cytometry. Optimal antibody dilution should be determined by titration; recommended range is 1:100 – 1:200 for flow cytometry, and for immunohistochemistry with avidin-biotinylated Horseradish peroxidase complex (ABC) as detection reagent, and 1:100 – 1:1000 for immunoblotting applications.

Storage

The antibody is shipped at ambient temperature and may be stored at +4°C. For prolonged storage prepare appropriate aliquots and store at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance or the concentration of the product.

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. It may contain hazardous ingredients. Please refer to the Safety Data Sheets (SDS) for additional information and proper handling procedures. Dispose product remainders according to local regulations. This datasheet is as accurate as reasonably achievable, but Exalpha Biologicals accepts no liability for any inaccuracies or omissions in this information.

References

1. Ramaekers, F. C., Puts, J. J., Moesker, O., Kant, A., Huysmans, A., Haag, D., Jap, P. H., Herman, C. J., and Vooijs, G. P. (1983). Antibodies to intermediate filament proteins in the immunohistochemical identification of Human tumours: an overview, *Histochem J* 15,

691-713. 2. Puts, J. J., Moesker, O., Kenemans, P., Vooijs, G. P., and Ramaekers, F. C. (1985). Expression of Cytokeratins in early neoplastic epithelial lesions of the uterine cervix, *Int J Gynecol Pathol* 4, 300-13. 3. Broers, J. L., Carney, D. N., Klein Rot, M., Schaart, G., Lane, E. B., Vooijs, G. P., and Ramaekers, F. C. (1986). Intermediate filament proteins in classic and variant types of small cell lung carcinoma cell lines: a biochemical and immunochemical analysis using a panel of monoclonal and polyclonal antibodies, *J Cell Sci* 83, 37-60. 4. Ramaekers, F., Huysmans, A., Schaart, G., Moesker, O., and Vooijs, P. (1987). Tissue distribution of Keratin 7 as monitored by a monoclonal antibody, *Exp Cell Res* 170, 235-49. 5. Van Muijen, G. N., Warnaar, S. O., and Ponec, M. (1987). Differentiation-related changes of Cytokeratin expression in cultured Keratinocytes and in fetal, newborn, and adult epidermis, *Exp Cell Res* 171, 331-45. 6. Bijman, J.T., Wagener, D.J.T., van Rennes, H., Wessels, J.M.C., Ramaekers, F.C.S. and van den Broek, P. (1987). Modulation of placental alkaline phosphatase activity and Cytokeratins in Human HN-1 cells by butyrate, retinoic acid, Catecholamines and histamine, *Br. J. Cancer* 56, 127-32. 7. Van Erp, P.E.J., Rijzewijk, J.J., Boezeman, J.B.M., Leenders, J., de Mare, S., Schalkwijk, J., van de Kerkhof, P.C.M., Ramaekers, F.C.S. and Bauer, F.W. (1989). Flow Cytometric analysis of epidermal subpopulations from normal and psoriatic skin using monoclonal antibodies anti intermediate filaments, *Am J Pathol* 135, 865-70. 8. Ramaekers, F., van Niekerk, C., Poels, L., Schaafsma, E., Huijsmans, A., Robben, H., Schaart, G., and Vooijs, P. (1990). Use of monoclonal antibodies to Keratin 7 in the differential diagnosis of adenocarcinomas, *Am J Pathol* 136, 641-55. 9. Schaafsma, H. E., Ramaekers, F. C., van Muijen, G. N., Lane, E. B., Leigh, I. M., Robben, H., Huijsmans, A., Ooms, E. C., and Ruiter, D. J. (1990). Distribution of Cytokeratin polypeptides in Human transitional cell carcinomas, with special emphasis on changing expression patterns during tumor progression, *Am J Pathol* 136, 329-43. 10. Smedts, F., Ramaekers, F., Robben, H., Pruszczynski, M., van Muijen, G., Lane, B., Leigh, I., and Vooijs, P. (1990). Changing patterns of Keratin expression during progression of cervical intraepithelial neoplasia, *Am J Pathol* 136, 657-68. 11. Franssen, M.E.J., Boezeman, J.B.M., van de Kerkhof, P.C.M. and van Erp, P.E.J. (2003). Monitoring hyperproliferative disorders in Human skin: flow cytometry of changing Cytokeratin expression, *Cytometry Part B (Clinical Cytometry)* 57B, 32-39.

Protein Reference(s)

Database Name: UniProt

Accession Number: P13645

Safety Datasheet(s) for this product:

NM_Sodium Azide