



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Mouse anti Cytokeratin 14 / Keratin K14

 nordicmubio.com/products/mouse-anti-cytokeratin-14-keratin-k14/MUB0323P

Catalog number: **MUB0323P**

Clone	RCK107
Isotype	IgG1
Product Type	Primary Antibodies
Units	0.1 mg
Host	Mouse
Species Reactivity	Canine Human Rat Swine
Application	Flow Cytometry Immunocytochemistry Immunohistochemistry (frozen) Western Blotting

Background

Cytokeratins are a subfamily of intermediate filament proteins and are characterized by a remarkable biochemical diversity, represented in Human epithelial tissues by at least 20 different polypeptides. They range in molecular weight between 40 kDa and 68 kDa and isoelectric pH between 4.9 – 7.8. The individual Human Cytokeratins are numbered 1 to 20. The various epithelia in the Human body usually express Cytokeratins which are not only characteristic of the type of epithelium, but also related to the degree of maturation or differentiation within an epithelium. Cytokeratin subtype expression patterns are used to an increasing extent in the distinction of different types of epithelial malignancies. The Cytokeratin antibodies are not only of assistance in the differential diagnosis of tumors using immunohistochemistry on tissue sections, but are also a useful tool in cytopathology and flow cytometric assays.

Source

RCK107 is a Mouse monoclonal IgG1 antibody derived by fusion of SP2/0-Ag14 Mouse myeloma cells with spleen cells from a Mouse immunized with a cytoskeletal preparation of TR146 epithelial cells.

Product

Each vial contains 100 ul 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.

Formulation: Each vial contains 100 ul 1 mg/ml purified monoclonal antibody in PBS containing 0.09% sodium azide.

Specificity

RCK107 reacts exclusively with Cytokeratin 14 which is present in basal cell compartments of stratified and combined epithelia.

Applications

RCK107 is suitable for immunocytochemistry, immunohistochemistry on frozen sections, flow cytometry and immunoblotting. Optimal antibody dilution should be determined by titration; recommended range is 1:100 – 1:200 for flow cytometry, and for immunohistochemistry with avidin-biotinylated Horseradish peroxidase complex (ABC) as detection reagent, and 1:100 – 1:1000 for immunoblotting applications.

Storage

The antibody is shipped at ambient temperature and may be stored at +4°C. For prolonged storage prepare appropriate aliquots and store at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance or the concentration of the product.

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. It may contain hazardous ingredients. Please refer to the Safety Data Sheets (SDS) for additional information and proper handling procedures. Dispose product remainders according to local regulations. This datasheet is as accurate as reasonably achievable, but Exalpha Biologicals accepts no liability for any inaccuracies or omissions in this information.

References

1. Wetzels, R. H., Kuijpers, H. J., Lane, E. B., Leigh, I. M., Troyanovsky, S. M., Holland, R., van Haelst, U. J., and Ramaekers, F. C. (1991). Basal cell-specific and hyperproliferation-related Keratins in Human breast cancer, *Am J Pathol* 138, 751-63.
- 2.

Smedts, F., Ramaekers, F., Troyanovsky, S., Pruszczynski, M., Robben, H., Lane, B., Leigh, I., Plantema, F., and Vooijs, P. (1992). Basal-cell Keratins in cervical reserve cells and a comparison to their expression in cervical intraepithelial neoplasia, *Am J Pathol* 140, 601-12. 3. Wetzels, R. H., Schaafsma, H. E., Leigh, I. M., Lane, E. B., Troyanovsky, S. M., Wagenaar, S. S., Vooijs, G. P., and Ramaekers, F. C. (1992). Laminin and type VII collagen distribution in different types of Human lung carcinoma: correlation with expression of Keratins 14, 16, 17 and 18, *Histopathology* 20, 295-303. 4. Bauwens, L. J., De Groot, J. C., Ramaekers, F. C., Veldman, J. E., and Huizing, E. H. (1992). Expression of intermediate filament proteins in the adult Human vestibular labyrinth, *Ann Otol Rhinol Laryngol* 101, 479-86. 5. Smedts, F., Ramaekers, F., Troyanovsky, S., Pruszczynski, M., Link, M., Lane, B., Leigh, I., Schijf, C., and Vooijs, P. (1992). Keratin expression in cervical cancer, *Am J Pathol* 141, 497-511. 6. Vos, J. H., van den Ingh, T. S., de Neijs, M., van Mil, F. N., Ivanyi, D., and Ramaekers, F. C. (1992). Immunohistochemistry with Keratin monoclonal antibodies in canine tissues: urogenital tract, respiratory tract, (neuro-)endocrine tissues, choroid plexus and spinal cord, *J Vet Med* 39, 721-40. 7. Vos, J. H., van den Ingh, T. S., Ramaekers, F. C., Molenbeek, R. F., de Neijs, M., van Mil, F. N., and Ivanyi, D. (1993). The expression of Keratins, vimentin, neurofilament proteins, smooth muscle actin, neuron-specific enolase, and synaptophysin in tumors of the specific glands in the canine anal region, *Vet Pathol* 30, 352-61. 8. van Leenders, G., Dijkman, H., Hulsbergen-van de Kaa, C., Ruiten, D., and Schalken, J. (2000). Demonstration of intermediate cells during Human prostate epithelial differentiation in situ and in vitro using triple-staining confocal scanning microscopy, *Lab Invest* 80, 1251-8. 9. Spies I. Immunolocation of mitochondria-rich cells in epidermis of the common toad, *Bufo bufo* L. (1997) *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 118,285-91.

Protein Reference(s)

Database Name: UniProt

Accession Number: P02533

Safety Datasheet(s) for this product:

NM_ Sodium Azide



Figure 1. Indirect immunofluorescence staining of swine skin frozen tissue section with MUB0323P (RCK107; Mouse anti keratin 14). Dilution 1:200. Specific staining of the basal and parabasal cells of the epidermis. No reactivity in the connective tissues.