



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Rabbit anti Chicken IgG (heavy and light chains)

Catalogue number: **RACH/IgG(H+L)**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Rabbit
Species reactivity	Chicken
Application	Immunoprecipitation

Background

The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in chicken serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In immunoelectrophoresis and double radial immunodiffusion to identify the presence of IgG; as secondary antibody to precipitate the immunoglobulin in normal chicken serum; to prepare an immunoabsorbent for the production of Ig-free chicken serum, plasma or non- Ig preparations.

Source

Purified normal IgG isolated from pooled chicken serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Product

Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit serum. No foreign proteins added.

Applications

Immunoprecipitation. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells

Cross Reactivity

Inter-species cross-reactivity is a normal feature of antibodies to

Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

immunoglobulins, since Ig of different species frequently share antigenic determinants. The degree of crossreactivity depends in addition to the phylogenetic relationship also on the concentration of the reactants and the sensitivity of the assay format. Cross-reactivity of this antiserum has not been tested in detail, however in immunoelectrophoresis a reaction with pigeon IgG has been observed.

Specificity

Precipitating polyclonal Rabbit antiserum to Chicken IgG, heavy and light chains.

Storage

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days.

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.