



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Rabbit anti Goat Ig light chains kappa and lambda (free and bound)

Catalogue number: **RAG/L(SD+HD)**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Rabbit
Species reactivity	Goat
Application	Immunoprecipitation

### Background

As tested in immunoelectrophoresis and double radial immunodiffusion against pooled normal goat serum and purified IgG fractions and light chain preparations, the antiserum reacts with determinants exposed on the surface (SD) and the Fab portion of all intact immunoglobulins, with Fab or F(ab')<sub>2</sub> subunits, and with free light chains of both types. This antiserum is not kappa or lambda chain specific. It does not react with any non-Ig protein in goat serum, as tested by immunoelectrophoresis and double radial immunodiffusion. As precipitating antiserum in a variety of immunodiffusion techniques including immunoelectrophoresis and double and single radial immunodiffusion (Mancini, Ouchterlony). It has not been tested for use in nephelometry, ELISA or immunochemistry, but this does not exclude such use if proper controls are included.

### Source

Purified light chains isolated from polyclonal goat IgG. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled rabbit serum. No foreign proteins added.

### Applications

Immunoprecipitation. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. In single immunodiffusion, use 1% antiserum in agar gel to establish a standard curve.

### **Cross Reactivity**

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail, but a wide scale of reactivity with IgG of other mammalian species must be assumed. The degree of cross-reactivity depends in addition to the phylogenetic relationship also on the concentrations of the reactants and the sensitivity of the assay format.

### **Specificity**

Precipitating polyclonal Rabbit antiSerum to Goat light chains, free and bound.

### **Storage**

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water. Working dilutions are prepared by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product Lyophilized at +4°C—at least 10 years. Reconstituted at or below -20°C—3-5 years. Reconstituted at +4°C—7 days.

### **Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.