



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Rabbit anti Hamster Serum proteins

Catalogue number: **RAHa/TSP**

Clone	Polyclonal
Product Type	Primary Antibodies
Units	1 ml
Host	Rabbit
Species reactivity	Hamster
Application	Immunoprecipitation

### Background

In immunoelectrophoresis against pooled serum and enriched serum proteins fractions precipitation can be observed of not less than 10 individual proteins components. In precipitating techniques as immunoelectrophoresis and radial immunodiffusion (Ouchterlony) to identify the serum protein pattern, or the presence or absence of an individual component. To evaluate the purity of an isolated serum protein including immunoglobulins. Since immunoprecipitation depends on a correct antigen/antibody concentration ratio (zone of equivalence) in the gel medium, the protein analysis by immunoelectrophoresis of serum or any other biological fluid or protein fraction should include different proportions of the reactants. It is not possible to obtain an optimal protein pattern in a single analysis. The electroendosmosis effect of different types of agar on proteins with a different net charge can be used to optimize the resolution power of the test system. Agar Nordic Nr. 2 contains sufficient positively charged ions to optimize the resolution of the proteins in the beta-gamma regions, while the alpha regions will become more dense. Highly purified agar (Agar Nordic nr. 1) with low electroendosmosis favours the resolution of the proteins in the alpha regions, while the major components in the beta-tgamma region can still be identified.

### Source

Pooled whole hamster serum and partly purified serum fractions. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

**Applications**

Immunoprecipitation. Different bleedings of the immunized animals are pooled to obtain a broad spectrum balanced against the varying concentrations of the individual serum protein components.

**Cross Reactivity**

Inter-species cross reactivity is a normal feature of antibodies to animal proteins since homologous proteins of different species frequently share antigenic determinants. This antiSerum has not been adsorbed for such cross reactivity. Consequently it is not species-specific.

**Specificity**

Precipitating polyclonal Rabbit antiSerum to Hamster Serum proteins.

**Storage**

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffer 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days.

**Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUBio accepts no liability for any inaccuracies or omissions in this information.