



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 IgA IgM (heavy and light chains), conjugated with Biotin

Catalogue number: **RAM/Ig/Bio**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Rabbit
Species reactivity	Mouse
Application	Dot blot ELISA Immunoblotting Immunocytochemistry Immunohistochemistry (paraffin)

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

### Background

The reactivity of the antiserum is directed the major immunoglobulin isotypes of the mouse Ig system (classes, subclasses and L-chain types), including antibodies to common determinants, to class and subclass-specific determinants, and to the surface determinants of the common Fab portion, as tested by immunoelectrophoresis and double radial immunodiffusion using various antiserum concentrations against normal mouse plasma and serum. In immunocytochemical and immunohistochemical staining of immunoglobulins at the cellular and subcellular level of appropriately treated cell and tissue substrates; to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of mouse origin in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure immunoglobulins in mouse serum or other body fluids. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

### Source

Purified polyclonal IgG and pools of homogenous IgA and IgM isolated from mouse serum. Freund's complete adjuvant is used in

the first step of the immunization procedure.

#### **Product**

Biotin-coupled purified hyperimmune goat IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity.

#### **Applications**

ELISA, Immunocytochemistry, Immunohistochemistry (paraffin), Dot blot, Immunoblotting.

#### **Cross Reactivity**

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. The antiSerum has been tested for cross-reactivity in double radial immunodiffusion with the following results: Bovine + Duck - Horse + Rat ++ Cat + Goat + Human + Sheep + Chicken - Guinea Pig ++ Monkey + Swine + Dog + Hamster ++ pigeon - Turkey ± A negative cross-reaction in double radial immunodiffusion does not exclude some reaction in more sensitive techniques.

#### **Specificity**

Biotin-conjugated IgG fraction of polyclonal Goat antiSerum to Mouse IgG, IgA and IgM, heavy and light chains

#### **Storage**

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days

#### **Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.