



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



## Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (heavy and light chains), conjugated with FITC

Catalogue number: **RAM/IgG(H+L)/FITC**

|                    |   |
|--------------------|---|
| Clone              | Polyclonal  |
| Product Type       | Secondary Antibodies  |
| Units              | 2 ml  |
| Host               | Rabbit  |
| Species reactivity | Mouse   |
| Application        | ELISA<br>Immunocytochemistry<br>Immunofluorescence<br>Immunohistochemistry (frozen) |

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

### Background

The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in mouse serum, as tested by immunoelectrophoresis and double radial immunodiffusion. To identify and measure IgG, antigen or antibody, at the cellular and subcellular level by immunofluorescence staining of appropriately treated cell and tissue substrates, and to demonstrate circulating antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen or immune complex using a reference antibody of mouse origin in the middle layer of the indirect test procedure. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

### Source

Purified normal IgG isolated from pooled mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

FITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added

**Applications**

ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.

**Cross Reactivity**

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.

**Specificity**

Fluorescein isothiocyanate-conjugated IgG fraction of polyclonal Rabbit antiSerum to Mouse IgG, heavy and light chains

**Storage**

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 2 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days

**Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.