



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Rabbit anti Mouse IgG1 (subclass specific)

Catalogue number: **RAM/IgG1**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Rabbit
Species reactivity	Mouse
Application	Immunoprecipitation

### Background

The reactivity of the antiserum is directed to the subclass IgG1. It does not react with other subclasses of IgG, IgG/Fab fragments, IgM and IgA or any non-Ig protein in mouse serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgG1 in mouse serum and other body fluids or to determine its concentration. To prepare an immunoadsorbent for the purification of mouse IgG1 from serum or plasma.

### Source

Pools of purified homogenous IgG1 isolated from pooled mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure

### Product

Delipidated, heat inactivated, lyophilized whole antiserum No preservative added, as it may interfere with the antibody activity Total protein and IgG concentration in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.

### Applications

Immunoprecipitation. In immunoelectrophoresis use 2 µl or equivalent against 120 µl antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in a 3 mm diameter centre well and 2 µl serum samples (neat and diluted) in 2 mm diameter peripheral wells

### Cross Reactivity

This IgG fraction is not species-specific since inter-species cross-reactivity is a normal feature of antisera to immunoglobulins.

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

Cross-reactivity of this product has not been tested in detail

**Specificity**

Precipitating polyclonal Rabbit antiSerum to Mouse IgG1, subclass specific

**Storage**

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C.

Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day.

Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days

**Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.