



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Rabbit anti Mouse milk specific proteins

Catalogue number: **RAM/MSP**

Clone	Polyclonal
Product Type	Primary Antibodies
Units	1 ml
Host	Rabbit
Species reactivity	Mouse
Application	Immunoprecipitation

### Background

In immunoelectrophoresis against mouse milk precipitation of not less than 7 different proteins may be obtained. However the number of visible precipitin lines varies among individual specimens of mouse milk. The total concentration of protein, their proportion of the different protein components change markedly during the course of lactation. There are also strain differences in the protein composition, but differences among individual animals tend to be greater. A protein detected in a mouse secretion by this antiserum is of secretory origin. However an analogous or homologous protein may be present also in other kind of secretions (e.g. saliva). Not all precipitable proteins have been identified. Known proteins include free secretory component and lactoferrin. In immunoelectrophoresis to identify the milk protein pattern, to compare individual samples and to make comparisons with serum protein patterns or those of other secretions; to identify individual components, their possible origin and significant changes in concentration. To test the purity of an isolated milk protein..

### Source

Pooled mouse milk. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.

### Applications

Immunoprecipitation. In immunoelectrophoresis use 2 µl milk or equivalent against 120 µl antiserum. In double radial

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

immunodiffusion (Ouchterlony) use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl milk samples (neat and serially diluted in 2 mm diameter peripheral wells). Different bleedings of the immunized animals are pooled to obtain a broad spectrum balanced against the varying concentrations of the individual protein components in individual milk samples.

#### **Cross Reactivity**

Inter-species cross reactivity is a normal feature of antibodies to animal proteins since homologous proteins of different species frequently share antigenic determinants. This antiSerum has not been adsorbed for such cross reactivity. Consequently it is not species-specific.

#### **Specificity**

Precipitating polyclonal Rabbit antiSerum to Mouse milk-specific proteins

#### **Storage**

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days

#### **Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUBio accepts no liability for any inaccuracies or omissions in this information.