



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Rabbit anti Monkey IgG Fab, conjugated with TRITC

Catalogue number: **RAMon/Fab/TRITC**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Rabbit
Species reactivity	Monkey
Application	ELISA Immunocytochemistry Immunofluorescence Immunohistochemistry (frozen)

Distributors

For Purchasing Information,
please contact your local
distributor

[Find Distributor](#)

Background

The reactivity of the antiserum is directed to the Fab subunits of intact IgG, IgA IgM and other Ig classes of both light chain types, with their Fab or F(ab')₂ subunits, and with free light chains of kappa and lambda type. In immunoelectrophoresis and double radial immunodiffusion using various antiserum concentrations against pooled monkey serum a characteristic precipitin line is obtained, which shows a reaction of identity with the precipitin line of the purified Fab. Direct staining of fixed cell and tissue substrates; to demonstrate the intracellular presence of free or Ig-bound subunits of both kappa or lambda type. In general this conjugate is not recommended as direct or indirect screening reagent for If isotypes on surface membranes of vital lymphoid cells. The activity to the common Ig/Fab subunit may result in the staining of immunoglobulins bound to the Fc-receptors on non-lymphoid cells. Combinations of isotype-specific reagents should be used instead for this purpose. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.

Source

Purified Fab of normal IgG isolated from Rhesus monkey serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

Product

TRITC-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No

preservative added as it may interfere with the antibody activity.

Applications

ELISA, Immunocytochemistry, Immunohistochemistry (frozen), (In)direct immunofluorescence.

Cross Reactivity

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail, however in immunoelectrophoresis a reaction with Human has been observed.

Specificity

Tetramethylrhodamine isothiocyanate-conjugated IgG fraction of polyclonal Rabbit antiSerum to Monkey Fab of IgG

Storage

The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C. Prior to use, an aliquot is thawed slowly in the dark at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate. Lyophilized at +4°C—at least 10 years. Reconstituted at or below -20°C—3-5 years. Reconstituted at +4°C—7 days

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.