



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Sheep anti Guinea Pig IgM (Fc specific)

Catalogue number: **ShAGp/IgM(Fc)**

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Sheep
Species reactivity	Guinea Pig
Application	Immunoprecipitation

### Background

The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal guinea pig serum a single precipitin line is obtained which shows a reaction of identity with the precipitin line obtained with purified IgM. No precipitation reaction is obtained with purified IgG, IgA, and IgG/Fab fragments. In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgM in guinea pig serum and other body fluids or to determine its concentration. To prepare an immunoabsorbent for the purification of guinea pig IgM from serum or plasma.

### Source

Highly purified normal IgM isolated from pooled guinea pig serum. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added.

### Applications

Immunoprecipitation. In immunoelectrophoresis use 2  $\mu$ l or equivalent against 120  $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10  $\mu$ l in a 3 mm diameter center well and 2  $\mu$ l serum samples (neat and serially diluted) in 2 mm diameter peripheral wells.

### Cross Reactivity

The antiSerum does not cross-react with any other component of

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

the Guinea Pig Ig system. Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.

**Specificity**

Precipitating polyclonal Sheep antiSerum to Guinea Pig IgM, Fc specific.

**Storage**

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days.

**Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.