



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## Sheep anti Human Factor I

Catalogue number: **ShAHu/FI**

Clone	Polyclonal
Product Type	Primary Antibodies
Units	1 ml
Host	Sheep
Species reactivity	Human
Application	Immunoprecipitation

## Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

### Background

The reactivity of the antiserum is restricted to factor I. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against human EDTA plasma a single precipitin line is obtained and shows a reaction of full identity with the purified immunogen. In precipitating techniques as immunoelectrophoresis and single and double radial immunodiffusion (Mancini, Ouchterlony) to identify the presence of F I or to determine its concentration. The presence of non-precipitating antibodies has not been assayed. This does not exclude the use of the antiserum in non-precipitating antibody-binding techniques if proper controls are included.

### Source

Factor I is a serine protease composed of two disulphide-linked chains. The protein has a molecular weight of 88.000 and its mean concentration in human plasma is 35 µg/ml. It is involved in the regulation of the C3/C5 convertase of either pathway. Using C4-binding protein or factor H splits the a chains of C4b or C3b respectively, which includes their roles in the C3/C5 convertases. It has been isolated as a homogenous protein for use in the antiserum production. Freund's complete adjuvant is used in the first step of the immunization procedure.

### Product

Delipidated, heat inactivated, lyophilized, stable IgG fraction. No preservative added. Total protein and IgG concentration are comparable to those of pooled normal sheep serum. No foreign proteins added.

### Applications

Immunoprecipitation. In immunoelectrophoresis in agarose-plates use 2 µl human plasma or equivalent against 120 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl plasma samples

(neat and serially diluted) in 2 mm diameter peripheral wells. In electroimmunodiffusion (Laurell) use a concentration of 1% of the antiserum in the gel.

#### **Cross Reactivity**

The antiSerum does not cross-react with any other Human plasma proteins as tested in gel diffusion techniques. Inter-species cross-reactivity is a normal feature of antibodies to plasma proteins, since homologous proteins of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.

#### **Specificity**

Precipitating polyclonal Sheep antiSerum to Human factor I.

#### **Storage**

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days.

#### **Caution**

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.