

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Sheep anti Human IgG3 (subclass specific)

Catalogue number: ShAHu/IgG3

Clone	Polyclonal
Product Type	Secondary Antibodies
Units	1 ml
Host	Sheep
Species reactivity	Human
Application	Immunoprecipitation

Background

The activity of the antiserum is directed to human IgG3 subclass. It does not react with any non-lg protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In precipitating techniques as immunoelectrophoresis and double radial immunodiffusion to identify and measure the concentration of IgG3 subclass in serum or other body fluids. Human IgG constitutes about 75% of the total Immunoglobulins in the serum of normal adults. It consists of four subclasses which are distinguished in isotypic heterogeneity of the gamma chains, and in physicochemical, biological and effector properties. The term "subclass" designates the high degree of homology in amino acid sequences and antigenic structure. In normal adult serum, the approximate percentage composition of IgG with respect to its subclasses is: IgG1: 65%, IgG2: 23%, IgG3: 8% and IgG4: 4%. All four subclasses appear in both isotypes of the light chain. Normal serum thus contains eight different molecular forms of IgG, the absolute and relative distribution being genetically controlled and resulting from individual responses to the antigenicity of the environment. Antigenic determinants carrying isotypic subclass specificity can be situated on any of the three structurally very similar sub-regions o the gamma chains, known as domains. In IgG, some antigenic determinants are shared by two or three subclasses, but not by all four. Consequently, antibodies to such determinants will not be strictly subclass-specific. For the identification of human IgG3 in cells and tissues by immunofluorescence or immunocytochemical peroxidase techniques, or in body fluids by ELISA and Western blotting type procedures, NORDIMMUNE monoclonal antibody reagents to the individual subclasses are available.

Source

Purified homogenous IgG3 is isolated from human serum. Freund's

Distributors

For Purchasing Information, please contact your local distributor

Find Distributor

complete adjuvant is used in the first step of the immunization procedure.

Product

Delipidated, heat inactivated lyophilized stable whole serum. No preservative added, as it may interfere with the antibody activity. No foreign protein added. Total protein and IgG concentration in the antiserum are comparable to those of pooled sheep serum. No foreign proteins added.

Applications

Immunoprecipitation. Because of their characteristics and those of the subclass-specific antigenic determinants, the use of this antiserum requires special precautions. The specificity and reactivity depend largely on the conditions of the test system in which they are applied. The performance of NORDIC polyclonal subclass-specific antisera is guaranteed when they are used for the purposes described in RWP 5E/88, applying the special immunoprecipitation arrangement recommended for these purposes. In immunoelectrophoresis use 2 µl 1:8, 1:16 and 1:32 dilutions of pathological serum or equivalent against 60 µl antiserum. In double radial immunodiffusion use a rosette arrangement with 10 µl antiserum in 3 mm diameter center well and 2 µl serum samples (neat and serially diluted) in 2 mm diameter peripheral wells. Nordic also makes available a human standard serum with assigned values of the subclasses of IgG (NOR-01). The antigenic differences between subclasses are considerably less pronounced than those between classes or between light chain types. Subclass-specific determinants are only few in number and probably restricted in their structural exposure, which makes them relative weak immunogens. This explains why the titres of subclass-specific antisera are usually lower than those of class-specific antisera which react with many antigenic determinants commonly present on all molecules of the respective class.

Cross Reactivity

Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.

Specificity

Precipitating polyclonal Sheep antisera to Human IgG3, subclass specific.

Storage

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. It is reconstituted by adding 1 ml sterile distilled water. Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days.

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures

involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.

Home | Company Profile | Catalogue | Distributors | Contact |

Content: Nordic-MUbio BV - Copyright © 2015