



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Sheep anti Human secretory IgA

Catalogue number: **ShAHu/sIgA**

| | |
|--------------------|----------------------|
| Clone | Polyclonal |
| Product Type | Secondary Antibodies |
| Units | 1 ml |
| Host | Sheep |
| Species reactivity | Human |
| Application | Immunoprecipitation |

Distributors

For Purchasing Information, please contact your local distributor

[Find Distributor](#)

Background

The reactivity of the antiserum is restricted to the Fc part of the IgA molecule and to surface and hidden determinants of the secretory component as tested in immunoelectrophoresis and radial immunodiffusion. In immunoelectrophoresis using various antiserum concentrations against normal human milk two precipitin lines may be obtained, one representing IgA and the other the free secretory component. Against normal human serum only IgA is precipitated. No precipitation reaction is obtained with purified IgG, IgM, and IgG/Fab fragments. In precipitating techniques as immunoelectrophoresis and radial immunodiffusion to identify the presence of IgA in human serum, milk or other body fluids or to determine its concentration. To prepare an immunoabsorbent for the purification of human IgA from serum or exocrine secretions. Antisera to IgA do not discriminate between serum IgA (monomeric and dimeric) and higher molecular forms as secretory IgA. This antiserum is not intended for use in non-precipitating antibody-binding or other highly sensitive assays. This does not exclude such use if proper controls are include.

Source

Highly purified secretory IgA isolated from pooled human milk. Freund's complete adjuvant is used in the first step of the immunization procedure.

Product

Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal sheep serum. No foreign proteins added.

Applications

Immunoprecipitation.

Cross Reactivity

The antiSerum does not cross-react with any other component of the Human Ig system. Interspecies cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.

Specificity

Precipitating polyclonal Sheep antiSerum to Human secretory IgA.

Storage

The lyophilized antiserum is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Reconstitute the lyophilized antiserum by adding 1 ml sterile distilled water. Dilutions may be prepared by adding phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the antiserum. Diluted antiserum should be stored at +4°C, not refrozen, and preferably used the same day. Lyophilized at +4°C--at least 10 years. Reconstituted at or below -20°C--3-5 years. Reconstituted at +4°C--7 days.

Caution

This product is intended FOR RESEARCH USE ONLY, and FOR TESTS IN VITRO, not for use in diagnostic or therapeutic procedures involving humans or animals. This datasheet is as accurate as reasonably achievable, but Nordic-MUbio accepts no liability for any inaccuracies or omissions in this information.