



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 009-001-106

Human Collagen Type IV

Overview

Description:	Human Collagen Type IV - 009-001-106
Item No.:	009-001-106
Size:	500 µg
Applications:	SDS-PAGE, ELISA
Origin:	Human

Product Details

Background:	Collagen type IV is a type of collagen found primarily in the basal lamina. The type IV collagen C4 domain at the C-terminus is not removed in post-translational processing, and the fibers link head-to-head, rather than in parallel. Also, collagen type IV lacks the regular glycine in every third residue necessary for the tight, collagen helix. This makes the overall arrangement more sloppy with kinks. These two features cause collagen IV to form in a sheet, the form of the basal lamina. There are six human genes associated with collagen IV: COL4A1, COL4A2, COL4A3, COL4A4, COL4A5 and COL4A6.
Synonyms:	Type IV collagen, collagen 4, human collagen, Collagen alpha-4 (IV) chain
Species of Origin:	Human
Type:	Native Protein

Target Details

Gene Name:	COL4A1-COL4A6
Purity/Specificity:	This product has been prepared from human placenta and is chromatographically and immunologically pure. This product is free from other collagens, human serum proteins and non-collagen extracellular matrix proteins. This product reacts with anti-Collagen Type IV. Reaction with Rockland's anti-Collagen I, II, III, V or VI is negligible.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P08572• UniProtKB - P02462• UniProtKB - P29400• UniProtKB - Q01955

- [UniProtKB - P53420](#)
- [UniProtKB - Q14031](#)

Application Details

Tested Applications:	SDS-PAGE
Suggested Applications:	ELISA (Based on references)
Application Note:	Collagen type IV has been tested in SDS-Page and can be used as a control or standard in indirect trapping ELISA for quantitation of antigen using a standard curve, for immunoprecipitation and for other immunological assays as optimized by the user. Specific conditions should be optimized by user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000
IHC:	1:100-1:500
WB:	1:1000


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/ml by nanodrop at 205 nm
Buffer:	0.5 M Acetic Acid
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is six (6) months from date of receipt.


Images



ELISA

Knockdown of NORAD inhibits proliferation, inflammation and fibrosis in HG-induced HMCs. Levels of (G) Col. IV in HMCs measured via ELISA. **P<0.01 vs. NG; ##P<0.01 vs. HG + si-NC.


NORAD, non-coding RNA activated by DNA damage; NG, normal glucose; HG, high glucose; HMCs, human mesangial cells; si, small interfering RNA; NC, negative control Col. IV, type IV collagen. Figure 2. PMID: 34194552.



ELISA

NORAD reverses the inhibitory effects of miR-485 on HG-induced HMCs. (F) Contents of Col. IV, FN and PAI-1 in HG-induced HMCs measured via ELISA **P<0.01 vs. miR-NC; #P<0.05 vs. miR-485 mimics.


miR, microRNA; NORAD, non-coding RNA activated by DNA damage; HG, high glucose; HMCs, human mesangial cells; ov, overexpression vector; NC, negative control; Col. IV, type IV collagen; FN, fibronectin; PAI-1, plasminogen activator inhibitor 1. Figure 4. PMID: 34194552.



ELISA

NORAD knockdown inhibits proliferation, inflammation and fibrosis in HG-induced HMCs by regulating the miR-485/NRF1 axis. (G) Contents of Col. IV, FN and PAI-1 in HG-induced HMCs as determined via ELISA. **P<0.01 vs. si-NC; #P<0.05 vs. si-NORAD + inhibitor NC; &P<0.05 vs. si-NORAD + ov-NC.

miR, microRNA; NORAD, non-coding RNA activated by DNA damage; NRF1, nuclear respiratory factor 1; HG, high glucose; HMCs, human mesangial cells; ov, overexpression vector; si, small interfering RNA; NC, negative control; Col. IV, type IV collagen; FN, fibronectin; PAI-1, plasminogen activator inhibitor 1. Figure 6. PMID: 34194552.



SDS-PAGE

SDS-PAGE Results of Human Collagen Type IV. Lane 1: Human Collagen Type IV Reduced (5.0µg). Lane 2: Opal Prestained Molecular Weight Protein (p/n MB-210-0500). Lane 3: Human Collagen Type IV Non-Reduced (5.0µg). 4-20% Gel, Coomassie Stained. Observed MW: Type IV α 1, α 2, α 3 bands.

References

- Wang L et al. Knockdown of lncRNA NORAD inhibits the proliferation, inflammation and fibrosis of human mesangial cells under high glucose conditions by regulating the miR485/NRF1 axis. *Exp Ther Med.* (2021)

Disclaimer

No test method can provide total assurance that the hepatitis B virus, hepatitis C virus, human immunodeficiency virus, or any other infectious agents are absent. Thus, all blood products, including purified proteins derived from human blood sources, should be handled at Biosafety Level 2 as recommended by the CDC\NIH manual entitled Biosafety in Microbiological and Biomedical Laboratories for potentially infectious human serum, blood specimens or proteins derived from same. Source material for the human blood product supplied to your facility has been tested for the detection of HIV antibody, Hepatitis B surface antigen, antibody to Hepatitis C, HIV 1 antigen(s), antibody to HTLV - I/II, and syphilis by FDA guidelines. All units were found to be non-reactive/negative for these tests. All human blood source material is collected in FDA licensed centers and is tested with FDA approved test kits.

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.