



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 009-001-B92-0.1

IL-3 Recombinant Human Protein

Overview

Description:	IL-3 Recombinant Human Protein - 009-001-B92-0.1
Item No.:	009-001-B92-0.1
Size:	100 µg
Applications:	SDS-PAGE
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	Interleukin-3 (IL-3) is a cytokine produced by activated T-cells and mast cells that is active on a broad range of hematopoietic cells and in the nervous system and appears to be important in several chronic inflammatory diseases. Recombinant Human IL-3 produced in E. coli is a single, non-glycosylated polypeptide chain containing 133 amino acids and having a molecular mass of 15,000 Daltons.
Synonyms:	Interleukin-3 cytokine, IL-3, Multipotential colony-stimulating factor, Hematopoietic growth factor, P-cell-stimulating factor, Mast cell growth factor, MCGF
Species of Origin:	Human
Expressed in:	E. coli
Type:	Recombinant Protein
Low Endotoxin:	Yes

Target Details

Gene Name:	IL3
Purity/Specificity:	Purity was determined to be greater than 95% as determined by analysis by RP-HPLC and by reducing and non-reducing SDS-PAGE, against known standard.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P08700• NCBI - 4E6EEURP015

Application Details

Tested Applications:	SDS-PAGE
Application Note:	IL-3 protein has been tested by SDS-PAGE and is suitable as a control for polyclonal or monoclonal anti-IL-3 in immunological assays. Buffer formation: 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	Biological Activity: Human IL-3 is fully biologically active when compared to standard. The ED50 as determined by the dose-dependent stimulation of TF-1 cells is less than 2 ng/ml. Endotoxin Level: Measured by LAL is <1EU/ μ g.


Formulation

Physical State:	Lyophilized
Concentration:	0.1mg/ml by UV absorbance at 280 nm
Buffer:	See application note.
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	100 μ L
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL) . For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
Expiration:	Expiration date is six (6) months from date of receipt.

Images

**SDS-PAGE**

SDS-PAGE shows bands corresponding to IL-3 (1 μ g) in lane 1 (non-reduced) and lane 3 (reduced, arrowhead). Molecular weight estimation was made by comparison to pre-stained MW markers, lane 2.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.