

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for 009-001-GM8

# **PARP1 (N-term ZF1) Control Protein**

#### **Overview**

Description:	PARP1 (N-term ZF1) Control Protein - 009-001-GM8
Item No.:	009-001-GM8
Size:	25 μg
Applications:	SDS-PAGE, WB
Origin:	Human
Expressed in:	E. coli

#### **Product Details**

Background:	PARP1 is the primary member of the poly(ADP-ribose) polymerase family, whose function is to signal DNA damage (and to recruit repair proteins) by PARylation. PARP1 is also involved in multiple cell death pathways, including apoptosis, necroptosis, autophagy, and a relatively new pathway termed parthanatos. PARP1 can also promote tissue survival by shifting the balance of cell death programs between autophagy and necrosis. Clinical studies have shown vulnerability to PARP inhibitors in DNA repair defective cancers. PARP1-ZF is useful for researchers interested in cellular processes including DNA damage, transcriptional control, and stem cell identity research.
Synonyms:	Poly [ADP-ribose] polymerase 1, ADP-ribosyltransferase diphtheria toxin-like 1, ARTD1, NAD(+) ADP-ribosyltransferase 1, ADPRT 1, PPOL
Species of Origin:	Human
Expressed in:	E. coli

## **Target Details**

Type:

Gene Name:	PARP1
Purity/Specificity:	PARP1 (N-term ZF1) is an N-terminus His-Tag recombinant protein expressed in E.coli that corresponds to a fragment of the human PARP1 zinc finger domain. Analysis by SDS-PAGE and Coomassie staining resulted in ~13 kDa MW band and estimated purity ≥90%.
Relevant Links:	• UniProtKB - P09874

**Recombinant Protein** 

www.rockland.com Page 1 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

# **Application Details**

<b>Tested Applications:</b>	SDS-PAGE, WB
Application Note:	PARP1 (N-term ZF1) protein has been tested in SDS-Protein and western blot and is suitable as a control for immunological assays that use Anti-PARP1-ZF (RABBIT) Antibody (200-401-GM8). Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 13 kDa in size corresponding to PARP-1 by western blotting.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

### **Formulation**

Liquid (sterile filtered)
1.0 mg/ml by modified Lowry assay
0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
0.01% (w/v) Sodium Azide
10% (v/v) Glycerol

# **Shipping & Handling**

<b>Shipping Condition:</b>	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable 2 - 3 weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.


## **Images**


www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823





#### **SDS-PAGE**

SDS-PAGE results of PARP1 (N-term ZF1) Control Protein. Lane 1: reduced PARP1 N-Term ZF1 protein. Lane 2: Opal Prestained Molecular Weight Ladder (p/n MB-210-0500). Lane 3: non-reduced PARP1 N-Term ZF1 protein. Load: 1µg. 4-20% Lonza SDS-PAGE; Coomassie Stained; BioRad ChemiDoc Imaged.

#### **Western Blot**

Western Blot of recombinant PARP1 with rabbit anti-PARP1 (N-term ZF1) antibody. Lane 1: PARP1-Zinc Finger domain recombinant protein. Load: 0.05  $\mu$ g per lane. Primary antibody: PARP1 (N-term ZF1) antibody at  $1\mu$ g/mL for overnight at 4°C. Secondary antibody: HRP Gt-a-rabbit secondary antibody (p/n 611-103-122) at 1:40,000 for 30 min at RT. Block: MB-070 overnight at 4°C. Predicted/Observed size: 13 kDa for rPARP1 (N-term ZF1). Other band(s): none.

#### **Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3