

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 009-001-HB1S

TRAF2 protein-GST fusion Epitope

Overview

Description:	TRAF2 recombinant protein-GST fusion Epitope - 009-001-HB1S
Item No.:	009-001-HB1S
Size:	20 μg
Applications:	SDS-PAGE, WB
Origin:	Human
Expressed in:	Sf9 cells

Product Details

Background: TRAF2 (also known as TNF receptor-associated factor 2) belongs to the TNF receptor associated

factor (TRAF) protein family that mediates signal transduction from members of the TNF receptor superfamily (1). TRAF2 is involved in TNF-alpha-mediated activation of MAPK8/JNK and NF-kB pathway. TRAF2 interacts with TNF receptors and forms a heterodimeric complex with TRAF1. The TRAF1-TRAF2 complex interacts with IAPs to mediate the anti-apoptotic signals from TNF receptors. TRAF2 also interacts with TRADD, a TNF receptor associated apoptotic signal transducer, to ensure the recruitment of IAPs for the direct inhibition of caspase activation (2). TRAF2 Protein is ideal for investigators involved in Signaling Proteins, Adaptor Proteins,

Apoptosis/Autophagy, Cancer, Inflammation, JNK/SAPK Pathway, and NfkB Pathway research.

Synonyms: TRAP, TRAP3, MGC: 45012, TNF receptor-associated factor 2, E3 ubiquitin-protein ligase TRAF2,

Tumor necrosis factor type 2 receptor-associated protein 3

Species of Origin: Human

Expressed in: Sf9 cells

Type: Recombinant Protein

Target Details

Gene Name: TRAF2

Purity/Specificity: Recombinant full-length human TRAF2 was expressed by baculovirus in Sf9 insect cells using an

N-Terminal his epitope. The purity was determined to be >80% by densitometry.

Relevant Links: • NCBI - NM_021138

www.rockland.com Page 1 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

Application Details

Tested Applications:	SDS-PAGE, WB
Application Note:	TRAF2 Protein is stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol. TRAF2 Protein is suitable for use in Western Blot. Expect a band approximately ~89kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
WB:	User Optimized

Formulation

Physical State:	Liquid (sterile filtered)
Buffer:	See application note.

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store product at -70 °C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.
Expiration:	Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

www.rockland.com Page 3 of 3