



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for 009-001-U65-0020**  
**rHuman BAFF - Protein****Overview**

<b>Description:</b>	Human BAFF - Recombinant Protein - 009-001-U65-0020
<b>Item No.:</b>	009-001-U65-0020
<b>Size:</b>	20 µg
<b>Applications:</b>	SDS-PAGE, Cellular Assay
<b>Origin:</b>	Human

**Product Details**

<b>Background:</b>	Human B cell Activating Factor (BAFF), is a Type II member of the TNF superfamily. Although it is typically a transmembrane protein expressed on T cells, macrophages and dendritic cells, it can also be cleaved in the extracellular region to produce a soluble form detectable in the serum. BAFF is thought to be important for the survival and maturation of peripheral B cells. BAFF initiates signaling through three receptors with human BAFF having activity on mouse splenocytes. Recombinant human BAFF is a non-glycosylated protein containing a N terminal His tag, a thrombin cut site and 153 amino acids of the TNF-like domain. This protein has a total molecular weight of 18.5 kDa.
<b>Synonyms:</b>	THANK, B lymphocyte stimulator (BLyS), B-cell activating factor, TNF- and APOL-related leukocyte expressed ligand 1 (TALL-1), CD_antigen=CD257
<b>Species of Origin:</b>	Human
<b>Type:</b>	Recombinant Protein
<b>Low Endotoxin:</b>	Yes

**Target Details**

<b>Gene Name:</b>	TNFSF13B
<b>Purity/Specificity:</b>	BAFF purity was determined to be greater than 90% as determined by analysis by HpLC, UV-Spectroscopy at 280nm, and by reducing and non-reducing SDS-pAGE.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - Q9Y275</a></li></ul>

## Application Details

<b>Tested Applications:</b>	SDS-PAGE
<b>Suggested Applications:</b>	Cellular Assay (Based on references)
<b>Application Note:</b>	Human BAFF Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-BAFF in immunological assays.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>Other:</b>	Endotoxin Level: Measured by kinetic LAL analysis and is typically $\leq 1$ EU/ $\mu$ g protein. Biologic Activity: The activity is determined by a RPMI-8226 cell survival assay in the presence of 0.2 $\mu$ M DEX and is typically less than 30 ng/mL.


## Formulation

<b>Physical State:</b>	Lyophilized
<b>Buffer:</b>	0.01 M Sodium Phosphate, pH 7.5
<b>Preservative:</b>	None
<b>Stabilizer:</b>	None
<b>Reconstitution Volume:</b>	20 $\mu$ l (20-200 $\mu$ l)
<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)


## Shipping & Handling

<b>Shipping Condition:</b>	Ambient
<b>Storage Condition:</b>	Store vial at -20° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
<b>Expiration:</b>	Expiration date is six (6) months from date of receipt.

## Images


**SDS-PAGE**

SDS-PAGE of Human BAFF Recombinant Protein. Lane 1: Molecular weight marker. Lane 2: 1 µg Human BAFF in non-reducing conditions (-). Lane 3: Molecular weight marker. Lane 4: 1 µg Human BAFF in reducing conditions (+). Human BAFF has a predicted MW of 17 kDa.


**SDS-PAGE**

Bioactivity of Human BAFF Recombinant Protein. RPMI-8226 cells were cultured with 0 to 500 ng/mL Human BAFF in the presence of 0.2 uM DEX. Cell proliferation was measured after 94 hours and the linear portion of the curve was used to calculate the ED50. The ED50 of Human BAFF is 23-35 ng/mL. This value is comparable to the competitor sample.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.