

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 009-001-V52-0010 rHuman IL-13 Protein

Overview

Description:	Human Interleukin-13 Recombinant Protein - 009-001-V52-0010
Item No.:	009-001-V52-0010
Size:	10 µg
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	Interleukin-13 (IL-13) is an important cytokine secreted from Th2 cells. The functions attributed to IL-13 overlap significantly with those of IL-4 (induces IgE secretion from B cells and inhibits expression of inflammatory cytokines such as IL-1 β , TNF- α , IL-8 and IL-6), but differs from IL-4 in that IL-13 seems to link inflammatory responses of immune cells to the pathophysiological changes in the surrounding non-immune cells. The receptor subunits of IL-13 consist of IL-4R α , IL-13R α 1 and IL-13R α 2. Human and mouse IL-13 are cross-reactive. Recombinant human IL-13 is a non-glycosylated variant of native IL-13, containing 115 amino acids (additional Gln at position 113), with a molecular weight of 12.6 kDa.
Synonyms:	NC30
Species of Origin:	Human
Expressed in:	E. coli
Туре:	Recombinant Protein
Low Endotoxin:	Yes

Target Details

Gene Name:	IL13
Purity/Specificity:	Interleukin-13 purity was determined to be greater than 95% as determined by analysis by reducing and non-reducing SDS-pAGE.
Relevant Links:	• UniProtKB - P35225



www.rockland.com tech@rockland.com +1 484.791.3823

Application Details

Application Note:	Interleukin-13 Recombinant Protein is suitable as a control for polyclonal or monoclonal anti- Interleukin-13 in immunological assays.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	Endotoxin Level: Measured by kinetic LAL analysis and is typically \leq 1 EU/µg protein. Biologic Activity: The activity is determined by the dose dependent proliferation of TF-1 cells and is typically less than 1 ng/mL.

Formulation

Physical State:	Lyophilized
Buffer:	0.01 M Sodium Citrate, pH 3.0
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	10µl (10-100µl)
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
Expiration:	Expiration date is six (6) months from date of receipt.

Disclaimer



Order online now!

www.rockland.com tech@rockland.com +1 484.791.3823

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.