

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 009-001-V53-0010

rHuman IL-15 Protein

Overview

Description:	Human Interleukin-15 Recombinant Protein - 009-001-V53-0010
Item No.:	009-001-V53-0010
Size:	10 μg
Applications:	Cellular Assay
Origin:	Human
Expressed in:	E. coli

Product Details

Background: Interleukin 15 (IL-15) is a widely expressed pro-inflammatory cytokine related to IL-2. IL-15

promotes activation of T cells, neutrophils and macrophages and is critical to dendritic cell function in several model systems. IL-15 has been shown to play a role in several inflammatory disorders, including rheumatoid arthritis, psoriasis and pulmonary inflammatory diseases. Emerging data suggests that IL-15 may serve as a good therapeutic target, as there appears to be a beneficial effect of IL-15 neutralization in models of psoriasis and diabetes. Human IL-15 shows activity on mouse cells. Recombinant human IL-15 is a non-glycosylated protein,

containing 114 amino acids, with a molecular weight of 12.8 kDa.

Synonyms: IL-T

Species of Origin: Human

Expressed in: E. coli

Type: Recombinant Protein

Low Endotoxin: Yes

Target Details

Gene Name: IL15

Purity/Specificity: Interleukin-15 purity was determined to be greater than 98% as determined by HpLC, analysis

by UV-Spectroscopy at 280nm, and by reducing and non-reducing SDS-pAGE.

Relevant Links: • UniProtKB - P40933

www.rockland.com Page 1 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

Application Details

Suggested Applications:	Cellular Assay (Based on references)
Application Note:	Interleukin-15 Recombinant Protein has been tested by biological activity and is suitable as a control for polyclonal or monoclonal anti-Interleukin-15 in immunological assays. Lyophilized in 10 mM sodium bicarbonate, pH 8.5.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	Endotoxin Level: Measured by kinetic LAL analysis and is typically ≤ 1 EU/µg protein. Biologic Activity: The activity is determined by the dose-dependant stimulation of the proliferation of mouse CTLL-2 cells and is typically less than 0.5 ng/mL.

Formulation

Physical State:	Lyophilized
Concentration:	0.1mg/ml
Buffer:	See application note.
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	10μl (10-100μl)
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling


Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
Expiration:	Expiration date is six (6) months from date of receipt.

Images

www.rockland.com Page 2 of 3



www.rockland.com tech@rockland.com +1 484.791.3823



SDS-PAGE

Bioactivity of Human Interleukin-15 Recombinant Protein. Serial dilutions of Human IL-15 (starting at 1 ng/mL) were added to CTLL-2 with other needed cytokines and stimuli. After 65 hours, cell viability was measured and the linear portion of the curve was us used to calculate the ED50. The ED50 of Lot # 0110-100-86 Human IL-15 is between 0.13-0.19 ng/mL. This value is comparable to the typical expected range of < 0.5 ng/mL.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3