



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 009-001-V72-0100**rHuman LIF Protein****Overview**

Description:	Human Leukemia Inhibitory Factor Recombinant Protein - 009-001-V72-0100
Item No.:	009-001-V72-0100
Size:	100 µg
Applications:	Cellular Assay
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	Human Leukemia Inhibitory Factor (LIF) is member of the IL-6 family and is made by a variety of tissues. Human LIF signals through gp130 to activate STAT3 in human ESCs . In mouse ESCs, LIF promotes self-renewal and pluripotency, but in human cells there is no indication that LIF has the same function. Instead, FGF-basic (FGF-2) and Activin A are used to promote self-renewal and pluripotent potential. Recombinant human LIF is a non-glycosylated protein, containing 181 amino acids, with a molecular weight of 19.9 kDa.
Synonyms:	Differentiation-stimulating factor (D factor), Leukocyte Inhibitory Factor, Melanoma-derived LPL inhibitor (MLPLI)
Species of Origin:	Human
Expressed in:	E. coli
Type:	Recombinant Protein
Low Endotoxin:	Yes

Target Details

Gene Name:	LIF
Purity/Specificity:	Leukemia Inhibitory Factor purity was determined to be greater than 98% as determined by reducing and non-reducing SDS-pAGE.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P15018

Application Details

Suggested Applications:	Cellular Assay (Based on references)
Application Note:	Leukemia Inhibitory Factor Recombinant Protein has been tested by biological activity and is suitable as a control for polyclonal or monoclonal anti-Leukemia Inhibitory Factor in immunological assays.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	Endotoxin Level: Measured by kinetic LAL analysis and is typically ≤ 1 EU/ μ g protein. Biologic Activity: The activity is determined by its ability to induce proliferation of TF-1 cells and is typically less than 200pg/mL.


Formulation

Physical State:	Lyophilized
Concentration:	0.1 mg/ml
Buffer:	0.1% Trifluoroacetic acid
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	100 μ L
Reconstitution Buffer:	0.01M acetic acid

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at -20° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
Expiration:	Expiration date is six (6) months from date of receipt.

Images



SDS-PAGE

Bioactivity of Human Leukemia Inhibitory Factor Recombinant Protein. Serial dilutions of Human LIF (starting at 20 ng/mL) were added to TF1 cells. After 66 hours, cell proliferation was measured and the linear portion of the curve was used to calculate the ED50. The ED50 of Human LIF is between 0.05-0.07 ng/mL.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.