

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 009-001-V76-0010 rHuman CCL7 Protein

Overview

Description:	Human Monocyte Chemotactic Protein-3 (CCL7) Recombinant Protein - 009-001-V76-0010
Item No.:	009-001-V76-0010
Size:	10 μg
Applications:	SDS-PAGE, Cellular Assay
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	Monocyte Chemotactic Protein 3 (MCP-3), also called CCL7, is produced by macrophages and some tumor cell lines. MCP-3 signals through three different G protein-coupled receptors, CCR1, CCR2, and CCR3. CCL7 chemoattracts monocytes and can regulate macrophage function. Recombinant human MCP-3 is a non-glycosylated protein, containing 76 amino acids, with a molecular weight of 9 kDa.
Synonyms:	CCL7, MARC
Species of Origin:	Human
Expressed in:	E. coli
Туре:	Recombinant Protein
Low Endotoxin:	Yes

Target Details

Gene Name:	MCP-3
Purity/Specificity:	Monocyte Chemotactic protein-3 (CCL7) purity was determined to be greater than 98% as determined by HpLC, analysis by UV-Spectroscopy at 280nm, and by reducing and non-reducing SDS-pAGE.
Relevant Links:	UniProtKB - Q7Z7Q8

www.rockland.com Page 1 of 3



www.rockland.com tech@rockland.com +1 484.791.3823

Application Details

Tested Applications:	SDS-PAGE
Suggested Applications:	Cellular Assay (Based on references)
Application Note:	Monocyte Chemotactic Protein-3 Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-Monocyte Chemotactic Protein-3 in immunological assays.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	Endotoxin Level: Measured by kinetic LAL analysis and is typically ≤ 1 EU/µg protein. Biologic Activity: The activity is determined by the ability to chemoattract human PBMCs, human neutrophils or THP-1 cells and is typically 10-100 ng/mL.

Formulation

Physical State:	Lyophilized
Buffer:	0.1% Trifluoroacetic acid
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	10µl (10-100µl)
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
Expiration:	Expiration date is six (6) months from date of receipt.


Images

www.rockland.com Page 2 of 3






www.rockland.com tech@rockland.com +1 484.791.3823



SDS-PAGE

SDS-PAGE of Human Monocyte Chemotactic Protein-3 (CCL7) Recombinant Protein. Lane 1: 1 μ g Human MCP-3 in non-reducing conditions (-). Lane 2: 1 μ g Human MCP-3 in reducing conditions (+). Lane 3: Molecular weight marker. Human MCP-3 has a predicted MW of 9 kDa.

Human MCP-3 Induced Chemotaxis of Human Neutrophils



SDS-PAGE

Bioactivity of Human Monocyte Chemotactic Protein-3 (CCL7) Recombinant Protein. Triplicate samples of primary human neutrophils from three donors were allowed to migrate to Human MCP-3 (10, 100 and 1000 ng/mL). After 30 minutes, cells that migrated were counted using a luminescent substrate and displayed on the bar graph above. Significant levels of migration over basal were seen in response to Human MCP-3/CCL7 starting at 10 ng/mL.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3