



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for 009-001-W34-0050**  
**rHuman TRAIL Protein****Overview**

<b>Description:</b>	Human TRAIL Recombinant Protein - 009-001-W34-0050
<b>Item No.:</b>	009-001-W34-0050
<b>Size:</b>	50 µg
<b>Applications:</b>	SDS-PAGE, Cellular Assay
<b>Origin:</b>	Human
<b>Expressed in:</b>	E. coli

**Product Details**

<b>Background:</b>	TNF-Related Apoptosis-Inducing Ligand (TRAIL) is produced by a wide variety of cell types and is shown to be a cytotoxic protein that induces apoptosis in tumor cells through activation of the death receptors, DR4 and DR5. Human TRAIL is active on mouse cells. Recombinant human TRAIL is a non-glycosylated protein, containing the 168 amino acid extracellular portion of TRAIL that is homologous to TNF. It has a molecular weight of 19.5 kDa.
<b>Synonyms:</b>	Apo2 Ligand (Apo2L), TL2, TNF-related apoptosis-inducing ligand, CD253
<b>Species of Origin:</b>	Human
<b>Expressed in:</b>	E. coli
<b>Type:</b>	Recombinant Protein
<b>Low Endotoxin:</b>	Yes

**Target Details**

<b>Gene Name:</b>	TNFSF10
<b>Purity/Specificity:</b>	TRAIL purity was determined to be greater than 90% as determined by analysis by UV-Spectroscopy at 280nm and by reducing and non-reducing SDS-pAGE.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - P50591</a></li></ul>

## Application Details

<b>Tested Applications:</b>	SDS-PAGE
<b>Suggested Applications:</b>	Cellular Assay (Based on references)
<b>Application Note:</b>	TRAIL Recombinant Protein has been tested by SDS-PAGE and biological activity and is suitable as a control for polyclonal or monoclonal anti-TRAIL in immunological assays. Product is provided as lyophilized powder in 0.05M Sodium Chloride and 0.01M Sodium Phosphate, pH 7.5.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>Other:</b>	Endotoxin Level: Measured by kinetic LAL analysis and is typically $\leq 1$ EU/ $\mu$ g protein. Biologic Activity: The activity is determined by the ability to induce apoptotic cell death in TRAIL-sensitive RPMI-8226 cells and is typically 10-15 ng/mL.


## Formulation

<b>Physical State:</b>	Lyophilized
<b>Buffer:</b>	See application note.
<b>Preservative:</b>	None
<b>Stabilizer:</b>	None
<b>Reconstitution Volume:</b>	50 $\mu$ L
<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)

## Shipping & Handling


<b>Shipping Condition:</b>	Ambient
<b>Storage Condition:</b>	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
<b>Expiration:</b>	Expiration date is six (6) months from date of receipt.

## Images



**SDS-PAGE**

SDS-PAGE of Human TRAIL Recombinant Protein. Lane 1: 1 µg Human TRAIL in non-reducing conditions (-). Lane 2: 1 µg Human TRAIL in reducing conditions (+). Lane 3: Molecular weight marker. Human TRAIL has a predicted MW of 19 kDa.



**SDS-PAGE**

Bioactivity of Human TRAIL Recombinant Protein. RPMI-8226 cells were cultured with 0 to 500 ng/mL Human TRAIL. Cell proliferation was measured after 5 days and the linear portion of the curve was used to calculate the ED50. The ED50 of Human TRAIL is 10-15 ng/mL. This value is comparable to the competitor sample, but there is no typically accepted range for this assay.

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.