



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 009-0131

Human IgM (myeloma) Fc5u

Overview

Description:	Human IgM (myeloma) Fc5 μ Fragment - 009-0131
Item No.:	009-0131
Size:	1 mg
Applications:	SDS-PAGE
Origin:	Human

Product Details

Background:	Human IgM (myeloma) Fc5 μ fragment consists of only the μ (mu) chain of the Fc fragment. Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. IgM is formed from covalently linking 5 immunoglobulins together. Due to this large size, IgM is typically isolated to the serum.
Synonyms:	Human immunoglobulin M fragment; Human IgM Fc5 μ
Species of Origin:	Human
Format:	IgM
Type:	Native Protein

Target Details

Purity/Specificity:	Human IgM (myeloma) Fc5 μ fragment has been prepared from Human IgM myeloma protein by digestion with trypsin followed by column chromatography. Purity was assessed by SDS-PAGE and HPLC to be greater than 95%. A single precipitin arc was observed against anti-human IgM Fc5 μ and anti-human serum when assayed by immuno-electrophoresis at a concentration of 20 mg/ml. No reaction was observed against anti-Trypsin, anti-human IgG F(ab') ₂ , anti-human IgG F(c), anti-human Kappa or anti-human Lambda.
----------------------------	---

Application Details

Tested Applications:	SDS-PAGE
-----------------------------	----------

Application Note:	Human IgM (myeloma) Fc5 μ fragment has been tested in SDS-Page and can be used in Western Blotting and ELISA experiments as a control reagent.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	User Optimized
IHC:	User Optimized
WB:	User Optimized


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.1 M Tris Chloride, 0.5 M Sodium Chloride, pH 8.0
Preservative:	0.05% (w/v) Sodium Azide

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. Human IgM (myeloma) Fc5 μ fragment is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Expiration:	Expiration date is one (1) year from date of receipt.

Images

**SDS-PAGE**

SDS PAGE Results of Human IgM Fc5 μ . Lane 1: Human IgM Fc5 μ reduced [5 μ g]. Lane 2: Opal Prestained Molecular Weight Marker (p/n MB-210-0500). Lane 3: Human IgM Fc5 μ non-reduced [5 μ g]. 4-20% gel, Coomassie stained. Bands at expected MW.

Disclaimer

No test method can provide total assurance that the hepatitis B virus, hepatitis C virus, human immunodeficiency virus, or any other infectious agents are absent. Thus, all blood products, including purified proteins derived from human blood sources, should be handled at Biosafety Level 2 as recommended by the CDC\NIH manual entitled Biosafety in Microbiological and Biomedical Laboratories for potentially infectious human serum, blood specimens or proteins derived from same. Source material for the human blood product supplied to your facility has been tested for the detection of HIV antibody, Hepatitis B surface antigen, antibody to Hepatitis C, HIV 1 antigen(s), antibody to HTLV - I/II, and syphilis by FDA guidelines. All units were found to be non-reactive/negative for these tests. All human blood source material is collected in FDA licensed centers and is tested with FDA approved test kits.

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.