

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 009-0609

Human IgM (myeloma) Fab mu Biotin

Overview

Description:	Human IgM (myeloma) Fab μ Fragment Biotin Conjugated - 009-0609
Item No.:	009-0609
Size:	1 mg
Origin:	Human

Product Details

Background:	Human IgM Fab mu Biotin is ideal for investigators involved in serum protein component research. IgM is by far the physically largest antibody in the human circulatory system. It is the first antibody to appear in response to initial exposure to antigen. The spleen is the major site of specific IgM production. Distinct heavy chains differ in size and composition; alpha and gamma contain approximately 450 amino acids, while μ and ϵ have approximately 550 amino acids.
Synonyms:	Human IgM Fab μ fragment Biotin conjugated, Human myeloma IgM Fab μ fragment Biotin conjugation
Species of Origin:	Human
Conjugate:	Biotin
Format:	IgM
Type:	Native Protein

Target Details

Purity/Specificity: This product was prepared from normal serum by delipidation, salt fractionation, ion exchange

> chromatography followed by enzyme digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Human IgM, anti-Human IgG F(ab')2 and anti-Human Serum. No reaction was observed against anti-Human IgM Fc5µ.

Application Details

Application Note: Monovalent Fab fragments of affinity-purified, secondary antibodies are offered to cover (block)

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

the surface of immunoglobulins for double labeling primary antibodies from the same host species, or to block endogenous immunoglobulins in tissue sections or on cell surfaces. They can be used for these purposes because each Fab fragment has only a single antigen binding site. HUMAN IgM (myeloma) Fab μ fragment Biotin conjugated can be utilized as a control or standard reagent in Western Blotting and ELISA experiments. Specific conditions should be optimized by the user.

Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be
	listed below.

Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Disclaimer

www.rockland.com Page 2 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

No test method can provide total assurance that the hepatitis B virus, hepatitis C virus, human immunodeficiency virus, or any other infectious agents are absent. Thus, all blood products, including purified proteins derived from human blood sources, should be handled at Biosafety Level 2 as recommended by the CDC\NIH manual entitled Biosafety in Microbiological and Biomedical Laboratories for potentially infectious human serum, blood specimens or proteins derived from same. Source material for the human blood product supplied to your facility has been tested for the detection of HIV antibody, Hepatitis B surface antigen, antibody to Hepatitis C, HIV 1 antigen(s), antibody to HTLV - I/II, and syphilis by FDA guidelines. All units were found to be non-reactive/negative for these tests. All human blood source material is collected in FDA licensed centers and is tested with FDA approved test kits.

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3