



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet for 010-001-V20-0100

**rMouse INF gamma Protein****Overview**

<b>Description:</b>	Mouse Interferon gamma Recombinant Protein - 010-001-V20-0100
<b>Item No.:</b>	010-001-V20-0100
<b>Size:</b>	100 µg
<b>Applications:</b>	Cellular Assay
<b>Origin:</b>	Mouse

**Product Details**

<b>Background:</b>	Interferon gamma, IFN $\gamma$ , is a major immune-modulating molecule produced mainly by T cells and natural killer cells activated by antigens, mitogens or alloantigens. Most immune cells express IFN $\gamma$ receptors and respond to IFN $\gamma$ -induced signaling by up-regulating MHC class I expression. Human IFN $\gamma$ does not show cross-reactivity with mouse. Recombinant mouse IFN $\gamma$ is a non-glycosylated protein, containing 134 amino acids, with a molecular weight of 15.6 kDa.
<b>Synonyms:</b>	Type II interferon, T cell interferon, Immune Interferon, MAF
<b>Species of Origin:</b>	Mouse
<b>Type:</b>	Recombinant Protein
<b>Low Endotoxin:</b>	Yes

**Target Details**

<b>Gene Name:</b>	Ifng
<b>Purity/Specificity:</b>	Interferon gamma purity was determined to be greater than 95% as determined by HPLC, analysis by UV-Spectroscopy at 280nm, and by reducing and non-reducing SDS-pAGE.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - P01580</a></li></ul>

**Application Details**

<b>Suggested Applications:</b>	Cellular Assay (Based on references)
--------------------------------	--------------------------------------

<b>Application Note:</b>	Interferon-gamma Recombinant Protein has been tested by biological activity and is suitable as a control for polyclonal or monoclonal anti-Interferon-gamma in immunological assays. Product is provided as a dried powder, lyophilized from 10 mM sodium phosphate, 100 mM sodium chloride, pH 7.5.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>Other:</b>	Endotoxin Level: Measured by kinetic LAL analysis and is typically $\leq 1$ EU/ $\mu$ g protein. Biologic Activity: The activity is determined in a viral challenge assay using EMC virus on L929 cells and is typically less than 0.2 ng/mL.


## Formulation

<b>Physical State:</b>	Lyophilized
<b>Concentration:</b>	0.1 mg/ml
<b>Buffer:</b>	See application note.
<b>Preservative:</b>	None
<b>Stabilizer:</b>	None
<b>Reconstitution Volume:</b>	100 $\mu$ L
<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)

## Shipping & Handling

<b>Shipping Condition:</b>	Wet Ice
<b>Storage Condition:</b>	Store vial at $-20^{\circ}$ C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at $-20^{\circ}$ C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
<b>Expiration:</b>	Expiration date is six (6) months from date of receipt.

## Images



### SDS-PAGE

Bioactivity of Mouse Interferon-gamma Recombinant Protein. The specific activity, as determined in a viral challenge assay using EMC virus on L929 cells, is 1.15-2.3 x 10<sup>7</sup> Units/mg. The corresponding ED50 is 0.086-0.043 ng/ml. The typical expected range of less than 0.2 ng/ml, corresponding to a specific activity of greater than 5 x 10<sup>6</sup> units/mg.

### Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.