



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet for 100-101-228

## N-Acylmannosamine-1-Dehydrogenase Antibody

### Overview

<b>Description:</b>	Anti-N-Acylmannosamine-1-Dehydrogenase (Pseudomonas) (GOAT) Antibody - 100-101-228
<b>Item No.:</b>	100-101-228
<b>Size:</b>	2 mL
<b>Applications:</b>	WB
<b>Reactivity:</b>	Pseudomonas
<b>Host Species:</b>	Goat

### Product Details

<b>Background:</b>	N-acylmannosamine 1-dehydrogenase acts on acetyl-D-mannosamine and glycolyl-D-mannosamine. NAM-DH may be applicable to the quantitative determination of sialic acid in serum.
<b>Synonyms:</b>	goat anti-N-Acylmannosamine-1-Dehydrogenase Antibody, N-acylmannosamine 1-dehydrogenase, NAM-DH
<b>Host Species:</b>	Goat
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	Antiserum

### Target Details

<b>Gene Name:</b>	NAM-DH
<b>Reactivity:</b>	Pseudomonas
<b>Immunogen Type:</b>	Recombinant Protein
<b>Immunogen:</b>	N-Acyl Mannosamine-1-Dehydrogenase [Recombinant Sequence from Pseudomonas expressed in E.coli ]
<b>Purity/Specificity:</b>	This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-goat serum, purified and partially purified N-Acyl Mannosamine-1-Dehydrogenase [E.coli Recombinant]. Cross reactivity against N-Acyl Mannosamine-1-Dehydrogenase from other sources is unknown.

<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - P22441</a></li><li>• <a href="#">NCBI - P22441.3</a></li><li>• <a href="#">GeneID - 7614553</a></li></ul>
------------------------	---

## Application Details

<b>Tested Applications:</b>	WB
<b>Application Note:</b>	Anti-N-Acyl Mannosamine-1-Dehydrogenase has been by western blot and is suitable to be assayed against 1.0 ug of N-Acyl Mannosamine-1-Dehydrogenase [E.coli Recombinant] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) code #605-4302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:5,000 to 1:20,000 of the reconstitution concentration is suggested for this product.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	1:5,000 - 1:20,000
<b>IP:</b>	1:100
<b>WB:</b>	1:500 - 1:5,000

## Formulation


<b>Physical State:</b>	Lyophilized
<b>Concentration:</b>	90 mg/mL by Refractometry
<b>Buffer:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.01% (w/v) Sodium Azide
<b>Stabilizer:</b>	None
<b>Reconstitution Volume:</b>	2.0 mL
<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)

## Shipping & Handling

<b>Shipping Condition:</b>	Ambient
----------------------------	---------

<b>Storage Condition:</b>	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
<b>Expiration:</b>	Expiration date is one (1) year from date of receipt.

## Images



### Western Blot

Western Blot Results of Goat Anti-N-Acylmannosamine-1-Dehydrogenase (Pseudomonas) Antibody.

Lane 1: SuperSignal Molecular Weight.

Lane 2: N-Acylmannosamine-1-Dehydrogenase.

Load: 0.05µg.

Primary Antibody: Goat Anti-N-Acylmannosamine-1-Dehydrogenase at 1:1000 overnight at 2-8°C.

Secondary Antibody: HRP Streptavidin (p/n S000-03) at 1:40,000 at RT for 30mins.

Blocking: Universal blocking buffer (p/n MB-073) for 30mins at RT.

Predicted/Observed MW: ~27.4kDa/~30kDa.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.