



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for 100-401-225****Dab1 Antibody****Overview**

<b>Description:</b>	Anti-Dab1 (400-555) (RABBIT) Antibody - 100-401-225
<b>Item No.:</b>	100-401-225
<b>Size:</b>	100 µL
<b>Applications:</b>	IHC, WB, Biochemical Assay, IF, Multiplex
<b>Reactivity:</b>	Mouse
<b>Host Species:</b>	Rabbit

**Product Details**

<b>Background:</b>	Anti-Dab1 Antibody recognizes Dab1 that is a phosphoprotein encoded by the mouse gene dab and is related to the Drosophila gene 'disabled'. Mutations in the mouse dab gene may result in the 'scrambler' and 'yotari' phenotypes. Dab1 binds to non-receptor tyrosine kinases and plays an important role in brain development. Dab1 is expressed in neuronal populations exposed to reelin, and it functions as a signaling molecule that regulates cell positioning in the developing brain. Cloning of human Dab1 and sequence determinations suggest a 96% identity to the mouse sequence. Dab1 binds to cytoplasmic regions of very low density lipoprotein receptors (VLDLR), apolipoprotein E receptor-2 (ApoER2) and the Amyloid Precursor Protein (APP) family of proteins. Dab1 accumulates in ectopic neurons from mice lacking Reelin or both VLDLR and ApoER2. In humans, Dab1 has been mapped to lp32-p312. This region shows homology of synteny with the segment of mouse chromosome 4 containing Dab1.
<b>Synonyms:</b>	rabbit anti-Dab 1 antibody, Disabled homolog 1 antibody, Disabled homolog 1 Drosophila antibody, Scm antibody, Scr antibody, Scrambler antibody, Yot antibody, Yotari antibody
<b>Host Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	Antiserum

**Target Details**

<b>Gene Name:</b>	Dab1
<b>Reactivity:</b>	Mouse

<b>Immunogen Type:</b>	Conjugated Peptide
<b>Immunogen:</b>	This whole rabbit serum was prepared by repeated immunizations with a synthetic peptide corresponding to the C-terminal region of murine Dab1 at amino acids 400-555.
<b>Purity/Specificity:</b>	Dab-1 whole rabbit antiserum was prepared by delipidation and defibrination followed by the addition of buffer salts and preservative. This antibody is directed against Dab1 from mouse. Cross-reactivity with other species has not been determined. No reaction occurs with human or mouse Dab2.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - P97318</a></li><li>• <a href="#">NCBI - NP_034144.1</a></li><li>• <a href="#">GeneID - 13131</a></li></ul>

## Application Details

<b>Tested Applications:</b>	IHC, WB
<b>Suggested Applications:</b>	Biochemical Assay, IF, Multiplex (Based on references)
<b>Application Note:</b>	Anti-Dab1 antibody has been tested by western blot and immunohistochemistry and is suitable for the detection of Dab1 by immunoprecipitation. Specific conditions for reactivity should be optimized by the end user. Expect a band at 80 kDa corresponding to Dab1 in the appropriate tissue extract or cell lysate. For western blotting block the blot using 5% BLOTTO for 1 h at room temperature and followed by incubation with the primary antibody diluted in 1% BLOTTO in TTBS for 1 h at room temperature. For immunoprecipitation use 1 µl of antiserum per 500 µg of brain lysate. Perform immunoprecipitation at 4°C for 2 h. For immunoprecipitation buffer lysates with 50 mM Tris-Cl, pH 7.4, supplemented with 150 mM sodium chloride, 1% (v/v) NP-40, 10 µg/ml aprotinin and 10 µg/ml leupeptin.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	1:10,000 - 1:50,000
<b>IF:</b>	User Optimized
<b>IHC:</b>	1:5,000
<b>IP:</b>	1 µl per 500 µg
<b>WB:</b>	1:5,000

## Formulation

<b>Physical State:</b>	Liquid (sterile filtered)
<b>Concentration:</b>	75 mg/mL by Refractometry

**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

---

**Preservative:** 0.01% (w/v) Sodium Azide

---

**Stabilizer:** None

---

## Shipping & Handling

**Shipping Condition:** Dry Ice

---


**Storage Condition:** Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

---


**Expiration:** Expiration date is one (1) year from date of receipt.

---

## Images


**A**

**Western Blot**

Coapplication of GST-RAP does not prevent the E2-induced distal dendritic HCN1 enrichment in vitro. A, Western blots showing phosphorylated Dab1 (pDab1, upper bands, detected by anti-phosphotyrosine antibody 4G10) and total Dab1 (lower bands) in "paired" slice cultures that were treated for 24 h (DIV10–11) with either GST-RAP or GST (10 μg/ml each) and were exposed to Reelin-conditioned medium for 30 min before harvesting. GAPDH was used for loading control. B, Quantitative analysis of Dab1 and pDab1 levels (relative to GAPDH) revealed that Reelin-induced pDab1 was reduced in the GST-RAP-treated slices, while total Dab1 was not significantly different compared to the GST-treated controls. Thus, pDab1/Dab1 was reduced to 72% ± 6% of control levels after 24-h GST-RAP treatment ( $p < 0.01$ ;  $n = 15$ ). C, D, E2 (+GST)-treatment (pink) of cultures for 6 d (DIV5–11) caused a significant HCN1 accumulation in segment 5 (C) and a significantly increased slope (D) compared with controls (GST, black) that was not efficiently reduced, if GST-RAP (10 μg/ml) was coapplied (orange;  $n = 17$ , each group). E–G, Representative photographs from a culture "triple," of which one culture served as a vehicle (GST)-treated control (E), while the others were treated with either E2 + GST (F) or E2 + GST-RAP (G). Note that HCN1 is accumulated at the hippocampal fissure (asterisks) at all conditions, but if E2 was present (F, G), less HCN1 immunosignal is visible in stratum radiatum (sr, arrows), indicating (relative) HCN1 enrichment in stratum lacunosum-moleculare (slm). Scale bars: 80 μm (E–G). Dashed lines demarcate the border of stratum pyramidale (sp), ml, molecular layer. Figure provided by CiteAb. Source: Eneuro, PMID: 30406178.




### Immunohistochemistry

Immunohistochemical staining after cryofixation and sectioning of mouse brain tissue using (A) a 1:5,000 dilution of anti-Dab-1 and (B) 1:5,000 dilution of pre-immune serum followed by processing with HRP Goat anti-Rabbit IgG [H&L] and chromogenic substrate.



### Western Blot

Analysis of GFP-Dab1 and endogenous Dab1 levels in transfected retinal cells and retinal tissue. Western blot analysis of whole cell lysates prepared from primary retinal cultures transfected with GFP-Dab1-E (lane 1), -L (lane 2), control untransfected retinal cultures (lane 3), and retinal tissue at ED5 (lane 4), ED7 (lane 5) and ED10 (lane 6). Proteins were electrophoresed through an SDS-8% polyacrylamide gel and transferred to nitrocellulose. The membrane was immuno-stained with Rockland Immunochemical's anti-Dab1 antibody at 1:5000, which recognizes both GFP-Dab1 (indicated by asterisks) and endogenous forms of Dab1 (indicated by a line). See Katval et al (2007) for additional details.



### Western Blot

Rockland Immunochemical's anti-Dab1 is shown to detect Dab1 present in wt mouse brain extracts (lane 1). No staining is noted in similar extracts from a dab knock-out mouse (lane 2). Detection of an 80 kDa band (arrowhead) occurs using a 1:5,000 dilution of the antibody in 1% milk in TTBS for 1 h at room temperature followed by a 1:5,000 dilution of HRP Goat-a-Rabbit with ECL visualization. Film exposure was ~1'. Other detection systems will yield similar results.

## References

- Wang G et al. The ZSWIM8 ubiquitin ligase regulates neurodevelopment by guarding the protein quality of intrinsically disordered Dab1. *Cereb Cortex*. (2023)
- Reyes, RV et al. The E3 Ubiquitin Ligase CRL5 Regulates Dentate Gyrus Morphogenesis, Adult Neurogenesis, and Animal Behavior. *Frontiers in Neuroscience* (2022)
- Kleene R et al. Serine 1283 in extracellular matrix glycoprotein Reelin is crucial for Reelin's function in brain development. *bioRxiv Preprint* (2022)
- Fairchild et al. RBX2 maintains final retinal cell position in a DAB1-dependent and -independent fashion. *Development* (2018)
- Meseke et al. Distal Dendritic Enrichment of HCN1 Channels in Hippocampal CA1 Is Promoted by Estrogen, but Does Not Require Reelin. *eNeuro* (2018)
- O'Brian B et al. Localization of the paranodal protein Caspr in the mammalian retina. *Mol Vis* (2010)
- de la Maza MP, Olivares D, Hirsch S, et al. Weight increase and overweight are associated with DNA oxidative damage in skeletal muscle. *Clin Nutr*. (2006)
- Bramblett DE et al. The transcription factor Bhlhb4 is required for rod bipolar cell maturation. *Neuron* (2004)
- Hanzlicek BW et al. Probing inner retinal circuits in the rod pathway: a comparison of c-fos activation in mutant mice. *Vis Neurosci*. (2004)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.