

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



www.rockland.com tech@rockland.com +1 484.791.3823

Datasheet for 100-401-J04 Eg5 Antibody

Overview

Description:	Anti-Eg5 (RABBIT) Antibody - 100-401-J04
Item No.:	100-401-J04
Size:	100 μL
Applications:	WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Eg5 (also known as kinesin-5 and KIF11) is a homo-tetramer which cross-links anti-parallel microtubules in the mitotic spindle to maintain spindle bipolarity. Eg5, a member of the Kinesin-5 subclass of kinesins, is a plus-end-directed tetrameric kinesin-family protein that influences the assembly and organization of the mitotic spindle, a self-assembled and dynamic microtubule-based structure that orchestrates chromosome segregation in dividing cells. Eg5 action is essential: when it is depleted from the cytoplasm of meiotically-mature Xenopus laevis eggs, abnormal monopolar spindles form, preventing successful division. Eg5 is expressed in all cells during mitosis and in post-mitotic neurons during development. In developing neurons pharmacological inhibition and siRNA knockdown of Eg5 results in longer axons, more branches, fewer bouts of axon retraction and the inability of growth cones to turn on contact with repulsive substrates. In migratory neurons, inhibition of Eg5 causes neurons to migrate in a random pattern and form shorter leading processes. In adult neurons, Eg5 has a similar effect on inhibiting the rate of short microtubule transport so pharmacological inhibition of adult Eg5 (i.e Monastrol) may be a potential therapeutic tool for the augmentation of adult axon regeneration.
Synonyms:	rabbit anti-Eg5 antibody, Kinesin-like protein KIF11, Kinesin-5, KIF11, Kinesin-related motor protein Eg5, Thyroid receptor-interacting protein 5
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	Antiserum

Target Details

ROCKLAND

www.rockland.com tech@rockland.com +1 484.791.3823

Gene Name:	KIF11
Reactivity:	Human
Immunogen Type:	Recombinant Protein
Immunogen:	Eg5 was prepared from whole rabbit serum produced by repeated immunizations with a truncated Eg5 construct expressed in E. coli corresponding to human Eg5 protein.
Purity/Specificity:	Anti-Eg5 is directed against the human Eg5 protein. The product was prepared from monospecific antiserum by delipidation and defibrination. A BLAST analysis was used to suggest reactivity with human. Cross-reactivity with Eg5 from other sources have not been determined.
Relevant Links:	 UniProtKB - P52732 NCBI - NP_004512.1 GeneID - 3799

Application Details

Tested Applications:	WB
Application Note:	Eg5 antibody has been tested by western blot. For western blots expect a band of approximately 72 kDa in size corresponding to truncated kinesin-1 protein. Specific conditions for reactivity should be optimized by the end user. This antibody is suitable for use in ELISA.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000
IHC:	User Optimized
WB:	1:1000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	46 mg/ml by Refractometry
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling



www.rockland.com tech@rockland.com +1 484.791.3823

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Rabbit Anti-Eg-5 Antibody. Lane 1: E.coli cell lysate expressing histidine tagged protein. Lane 2: Mouse brain lysate. Lane 3: Recombinant truncated Eg-5. Lane 4: MW Markers. Load: 35μ g/lane for cell lysate, 50ng of recombinant protein. Primary antibody: Eg-5 antibody at 1:1000 for overnight at 4°C. Secondary antibody: HRP rabbit secondary antibody (p/n 611-103-122) at 1:40,000 for 60 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 120 kDa and 72 kDa for Eg5.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.