



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for 200-101-037

Glucose Oxidase Antibody

Overview

Description:	Anti-Glucose Oxidase (GOAT) Antibody - 200-101-037
Item No.:	200-101-037
Size:	100 µg
Applications:	WB
Reactivity:	Aspergillus niger
Host Species:	Goat

Product Details

Background:	Glucose Oxidase enzyme is widely applied for the determination of glucose in body fluids and in removing residual glucose or oxygen from foods and beverages. Furthermore, glucose oxidase-producing molds such as aspergillus and penicillium species are used for the biological production of gluconic acid.
Synonyms:	goat anti-Glucose Oxidase Antibody, Beta D Glucose Oxygen 1 Oxido Reductase antibody, Glucose oxidase (Precursor) antibody, Glucose Oxyhydrase antibody, GOD antibody
Host Species:	Goat
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	gox
Reactivity:	Aspergillus niger
Immunogen Type:	Native Protein
Immunogen:	Glucose Oxidase [Aspergillus niger]

Purity/Specificity: Anti-Glucose Oxidase Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Glucose Oxidase [*Aspergillus niger*]. Cross reactivity against Glucose Oxidase from other tissues and species may occur but have not been specifically determined.

Relevant Links:

- [UniProtKB - P13006](#)
- [NCBI - P13006.1](#)

Application Details

Tested Applications:	WB
Application Note:	Anti-Glucose Oxidase has been tested in western blot and is suitable for use in ELISA and immunohistochemistry. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:2,000 - 1:10,000
IHC:	User Optimized
WB:	1:1000 - 1:5000

Formulation



Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/ml by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store Glucose Oxidase Antibody at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Rockland Goat anti Glucose Oxidase antibody was used to detect Glucose Oxidase under reducing and non-reducing conditions. Reduced samples of purified Glucose Oxidase contained 4% BME and were boiled for 5 minutes. Samples of ~1 and 0.25 ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with Rockland Goat anti-Glucose Oxidase antibody (p/n 200-1137, lot 165) at 1:5K in MB-070, ON 4°C. Primary antibody was detected with Rockland Dylight 649 conjugated Donkey anti-Goat (p/n 605-743-125, lot 20834) at 1:10K 1.5 hr RT in MB-070 and imaged on the BioRad VersaDoc imaging system.

Western Blot

Western Blot of Goat anti-Glucose Oxidase antibody. Lane 1: Glucose Oxidase. Load: 50 ng per lane. Primary antibody: Glucose Oxidase antibody at 1:1,000 for overnight at 4°C. Secondary antibody: Peroxidase goat secondary antibody (p/n 605-4302) at 1:40,000 for 30 min at RT. Block: (p/n MB-070) for 30 min at RT. Predicted/Observed size: 65-70 kDa, 65-70 kDa for Glucose Oxidase. Other band(s): None.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.