

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com



www.rockland.com tech@rockland.com +1 484.791.3823

# Datasheet for 200-4336 Beta Galactosidase Antibody Peroxidase Conjugated

#### **Overview**

| Description:  | Anti-Beta Galactosidase (E. coli) (RABBIT) Antibody Peroxidase Conjugated (BULK ORDER) - 200-<br>4336 |
|---------------|---|
| Item No.:     | 200-4336  |
| Size:         | 10 mg   |
| Applications: | ELISA, WB   |
| Reactivity:   | b-GAL   |
| Host Species: | Rabbit  |

#### **Product Details**

| Background:   | Anti Beta Galactosidase Antibody recognizes the enzyme beta galactosidase, or $\beta$ -galactosidase, that is a component of assays used frequently in genetics, molecular biology (see X-gal) for a blue white screen, and other life sciences. IPTG induces production of $\beta$ -galactosidase by binding and inhibiting the lac repressor. Since it is highly expressed and accumulated in lysosomes in senescent cells, it is used as a senescence biomarker both in vivo and in vitro in qualitative and quantitative assays, despite its limitations. Anti-beta Galactosidase Antibody is ideal for investigators involved in enzyme research. |
|---------------|--|
| Synonyms:     | rabbit anti-Beta Galactosidase Antibody Peroxidase Conjugation, HRP conjugated rabbit anti-<br>Beta Galactosidase Antibody, rabbit anti-beta gal HRP conjugated antibody   |
| Host Species: | Rabbit   |
| Conjugate:    | Peroxidase (HRP)   |
| Clonality:    | Polyclonal   |
| Format:       | lgG  |

### **Target Details**

| Reactivity:     | b-GAL                       |
|-----------------|-----------------------------|
| Immunogen Type: | Native Protein              |
| Immunogen:      | Beta Galactosidase (E.coli) |



#### Order online now!

www.rockland.com tech@rockland.com +1 484.791.3823

| Purity/Specificity: | This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step<br>process which includes delipidation, salt fractionation and ion exchange chromatography<br>followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis<br>resulted in a single precipitin arc against anti-Peroxidase, and anti-Rabbit Serum as well as<br>purified and partially purified Beta Galactosidase [E.coli]. Cross reactivity against Beta<br>Galactosidase from other tissues and species may occur but have not been specifically<br>determined. Very low background staining has been reported in various assays. |
|---------------------|---|
| Relevant Links:     | • UniProtKB - P00722  |
|                     | • NCBI - NP_414878.1  |

#### **Application Details**

| Tested Applications: | ELISA, WB  |
|----------------------|--|
| Application Note:    | Anti-Beta Galactosidase Peroxidase Conjugated Antibody has been tested by ELISA and Western<br>blot and is suitable for dot blot, immunofluorescence microscopy, immunoprecipitation,<br>conjugation and most immunological methods requiring high titer and specificity. The antibody<br>recognizes both frozen tissue sections, paraffin embedded tissue and 4% paraformaldehyde<br>fixed tissue for most immunohistochemical analysis. A 1:5,000 dilution has been reported to be<br>successful for staining by immunoblot of beta-galactosidase fusion proteins after transfer using<br>a semi-dry transfer apparatus. A 1:1,500 dilution has been reported to detect beta-<br>galactosidase in adult rat spinal cord tissue after infection with helper-dependent adenovirus<br>expressing lacZ. In this particular experiment, tissue was perfused with 4% paraformaldehyde<br>and cryostat-cut ( $35 \mu$ m) to produce free-floating sections. A 1:5,000 dilution has been reported<br>to be successful for staining of brain sections from transgenic mice expressing nuclear beta-<br>galactosidase when assayed by immunofluorescence microscopy. A 1:5,000 dilution has been<br>reported for immunofluorescent staining of methanol fixed, devitellinized Drosophila embryos.<br>Although a wide range of conditions was reported to be effective, a 1:10,000 dilution was noted<br>to show no background and to be suitable for double labeling experiments. Optimal titers for<br>other applications should be determined by the researcher. |
| Assay Dilutions:     | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.  |
| ELISA:               | 1:10,000 - 1:50,000  |
| IHC:                 | 1:500 - 1:2,000  |
| WB:                  | 1:2,000 - 1:5,000  |

### Formulation

| Physical State: | Lyophilized                           |
|-----------------|---------------------------------------|
| Concentration:  | 10.0 mg/mL by UV absorbance at 280 nm |



www.rockland.com tech@rockland.com +1 484.791.3823

| Buffer:                | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2             |
|------------------------|--|
| Preservative:          | 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!               |
| Stabilizer:            | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Reconstitution Volume: | 1.0 mL   |
| Reconstitution Buffer: | Restore with deionized water (or equivalent)                           |

#### **Shipping & Handling**

| Shipping Condition: | Ambient   |
|---------------------|---|
| Storage Condition:  | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| Expiration:         | Expiration date is one (1) year from date of receipt.   |

#### Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.