



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for BSA-30****Bovine Serum Albumin 30% Solution****Overview**

<b>Description:</b>	Bovine Serum Albumin 30% Solution - BSA-30
<b>Item No.:</b>	BSA-30
<b>Size:</b>	500 mL
<b>Applications:</b>	Biochemical Assay, ELISA, Other
<b>Origin:</b>	Bovine

**Product Details**

<b>Background:</b>	Bovine Serum Albumin (BSA) is used for various biochemical applications including ELISA (Enzyme-Linked Immunosorbent Assay), high content screening assays, western blotting, and immunohistochemistry. BSA as a blocking reagent is particularly useful with casein-sensitive antibodies, such as phospho-specific antibodies. Sterile validated preparations may be used as a nutrient in cell and microbial culture. In restriction digests, specially prepared BSA is used to stabilize some enzymes during digestion of DNA and to prevent adhesion of the enzyme to reaction tubes and other vessels. Bovine Serum Albumin can also be used to determine the quantity of other proteins, by comparing an unknown quantity of protein to known amounts of BSA.
<b>Synonyms:</b>	BOVINE SERUM ALBUMIN, BSA, BSA Blocker, BSA Blocking, BSA30
<b>Species of Origin:</b>	Bovine

**Target Details**

<b>Purity/Specificity:</b>	BOVINE SERUM ALBUMIN 30% Solution contains a 30% solution of bovine serum albumin (Fraction V, Immunoglobulin and Protease Free) in saline with preservative and stabilizer. The product is ~30% protein as determined by Biuret assay. A clear yellow green to dark amber color is typical. This product is prepared from healthy adult animals and is processed from pooled lots of serum to ensure uniformity.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li><a href="#">BSA-30 SDS</a></li></ul>

**Application Details**

<b>Suggested Applications:</b>	Biochemical Assay, ELISA, Other (Based on references)
<b>Application Note:</b>	BOVINE SERUM ALBUMIN 30% Solution is suitable for use in protease sensitive assays such as RIA, EIA and nucleic acid hybridization, use as a stabilizing agent for proteins and enzymes, including dilute solutions of antibody, and use as a blocking agent to reduce non-specific binding.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	User Optimized
<b>FLISA:</b>	User Optimized
<b>IF:</b>	User Optimized
<b>IHC:</b>	User Optimized
<b>WB:</b>	User Optimized


## Formulation

<b>Physical State:</b>	Liquid (sterile filtered)
<b>Concentration:</b>	30% by dry weight
<b>Buffer:</b>	0.85% (w/v) Sodium Chloride
<b>Preservative:</b>	0.01% (w/v) Sodium Azide
<b>Stabilizer:</b>	0.3% (v/v) Octanoic Acid

## Shipping & Handling


<b>Shipping Condition:</b>	Wet Ice
<b>Storage Condition:</b>	Store Blocking Buffer at 4° C prior to opening. DO NOT FREEZE.
<b>Expiration:</b>	Expiration date is six (6) months from date of receipt.

## Images



### Immunofluorescence Microscopy

Internalization of basigin monoclonal antibodies (mAbs) in hCMEC/D3 cells. (a) Representative confocal images of selected basigin mAbs exposed to hCMEC/D3 cells for 10 and 30 min. Basigin mAbs in green and nuclei visualized by Hoechst in blue. The pictures are the maximum projection of the z-stack, and the XZ projection is below the pictures. Scale bar 30  $\mu$ m. (b) Quantification of intracellular spots using Cellomics Arrayscan after acid stripping and staining. The intensities are normalized to the negative control and plotted as the percentage increase in spot intensity per cell with  $\pm$  standard error of the mean (SEM). Secondary antibody diluted in PBS with 2% BSA (p/n BSA-30). Figure 6. PMID: 32884039.



### Surface Plasmon Resonance (SPR)

Surface plasmon resonance sensorgrams representing the binding of basigin extracellular domain to captured anti-basigin monoclonal antibodies. The fitted curves used for equilibrium dissociation constant (KD) calculations were obtained using the Langmuir 1:1 binding model in the Biacore S200 evaluation software and plotted in GraphPad Prism (colored in black). (a) bin A, (b) bin D, (c) bin AD, (d) bin B, and (e) bin C. At the end of each cycle, the sensor surface was regenerated in 3 M MgCl<sub>2</sub> for 30 s. 1  $\times$  HBS-P + with 1 mg/ml bovine serum albumin (BSA, p/n BSA-30) was used as running buffer for the kinetic analyses. mAb monoclonal antibody, RU response unit. Figure 3. PMID: 32884039.

### Bottle

Bovine Serum Albumin 30% Solution



## References

- Yang X et al. Qualitative immunoassay for the detection of anti-SARS-COV-2 spike antibody in human milk samples. *STAR Protoc.* (2022)
- Christensen SC et al. Characterization of basigin monoclonal antibodies for receptor-mediated drug delivery to the brain. *Sci Rep.* (2020)
- Ferdman J et al. Intra-seasonal antibody repertoire analysis of a subject immunized with an MF59®-adjuvanted pandemic 2009 H1N1 vaccine. *Vaccine.* (2019)
- Raymond DD et al. Influenza immunization elicits antibodies specific for an egg-adapted vaccine strain. *Nat Med.* (2016)
- Schmidt AG et al. Viral receptor-binding site antibodies with diverse germline origins. *Cell.* (2015)
- Keeler, GD et al. Comparison of microdialysis sampling perfusion fluid components on the foreign body reaction in rat subcutaneous tissue. *European Journal of Pharmaceutical Sciences : Official Journal of the European Federation for Pharmaceutical Sciences* (2014)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.