



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet for D120-00-0050**Donkey Serum Sterile****Overview**

Description:	Donkey Serum (Sterile) - D120-00-0050
Item No.:	D120-00-0050
Size:	50 mL
Applications:	IF, IHC, Multiplex
Origin:	Donkey

Product Details

Background:	Sterile Donkey Serum is used as a supplement to cell culture media. Donkey Serum provides a broad spectrum of macromolecules, carrier proteins for lipoid substances and trace elements, attachment and spreading factors, low molecular weight nutrients, and hormones and growth factors that promote cell growth and health. Be certain to maintain Good Cell Culture Practice, and maintain sterility of cultures that require media supplementation.
Synonyms:	sterile Donkey serum for cell culture, cell culture grade donkey serum, sterile serum from donkey
Species of Origin:	Donkey

Application Details

Suggested Applications:	IF, IHC, Multiplex (Based on references)
Application Note:	pH: normal Immunoelectrophoresis: normal Hemoglobin: normal IgG Concentration: normal
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

Tissue Data

Tissue Type:	Serum
Sex:	Mixed
Strain:	Donkey - Mixed


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	92mg/ml by Refractometry
Buffer:	None
Sterility:	Sterile
Preservative:	None
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store container at -20° C prior to opening. Avoid cycles of freezing and thawing. Use aseptic technique to maintain sterility when opening product.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Immunofluorescence Microscopy

Loss of Stx5S compromises ER–Golgi trafficking.

(a) Immunofluorescence microscopy of GALNT2 (green in merge) in primary human dermal fibroblasts of healthy donors (green, Ctrl) or Stx5M55V patients (orange, Stx5M55V) in the absence or presence of brefeldin A (BFA) for 6 min and washout for the indicated times.

Representative confocal micrographs are shown. Following quenching with 50 mM NH₄Cl in PBS, cells were permeabilized and blocked in 2% normal donkey serum (p/n D120-00-0050) and 0.1% saponin (permeabilization buffer) for 30 min at RT. Scale bars, 10 μm. DAPI in blue. Fig. 5. PMID: 34711829.



Bottle

Donkey Serum is used as a supplement to cell culture media. Donkey Serum provides a broad spectrum of macromolecules, carrier proteins for lipid substances and trace elements, attachment and spreading factors, low molecular weight nutrients, and hormones and growth factors that promote cell growth and health. Be certain to maintain Good Cell Culture Practice, and maintain sterility of cultures that require media supplementation.

References

- Linders, PTA et al. Congenital disorder of glycosylation caused by starting site-specific variant in syntaxin-5. *Nature Communications* (2021)
- Hilgen G et al. Subcellular distribution of connexin45 in OFF bipolar cells of the mouse retina. *J Comp Neurol.* (2011)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.