



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

**Datasheet for D204-00-1000****Goat Serum Non-Sterile****Overview**

<b>Description:</b>	Goat Serum (Non-Sterile) - D204-00-1000
<b>Item No.:</b>	D204-00-1000
<b>Size:</b>	1 L
<b>Applications:</b>	IF, Other
<b>Origin:</b>	Goat

**Product Details**

<b>Background:</b>	Normal Goat Serum can be used as a blocking agent to treat plastic surfaces, membrane or tissue after they have been sensitized with primary antibody or antigen. It provides an alternative to bovine serum albumin (BSA) and non-fat dry milk. It is effective in reducing nonspecific binding of proteins to reaction surfaces, thereby maximizing signal-to-noise ratio. This blocking agent is recommended for use in immunoassays where the primary antibody was produced in goat, as a source of non-specific serum protein or on tissue for immunohistochemical applications. Normal Goat Serum is used as a component of bioassays, immunoassays or enzyme assays. Normal Goat Serum provides a broad spectrum of macromolecules, carrier proteins for lipid substances and trace elements, attachment and spreading factors, low molecular weight nutrients, and hormones and growth factors that promote cell growth and health. Normal Goat Serum is also available as a sterile preparation for use as a cell culture supplement. Normal Goat Serum is ideal for investigators in Cancer, Immunology and Cell Biology research.
<b>Synonyms:</b>	normal goat serum, NGS, blocking serum, goat serum for bioassay, control goat serum, non-sterile serum from goat, non-sterile goat serum
<b>Species of Origin:</b>	Goat

**Target Details**

<b>Purity/Specificity:</b>	Goat serum was obtained from healthy non-immunized goats.
----------------------------	---

**Application Details**

<b>Suggested Applications:</b>	IF, Other (Based on references)
--------------------------------	---------------------------------

---

<b>Application Note:</b>	pH: normal
	Immunoelectrophoresis: normal
	Hemoglobin: normal
	IgG Concentration: normal

---

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

---

<b>ELISA:</b>	User Optimized
<b>FC:</b>	User Optimized
<b>FLISA:</b>	User Optimized
<b>IF:</b>	User Optimized
<b>IHC:</b>	User Optimized
<b>IP:</b>	User Optimized
<b>WB:</b>	User Optimized

---

## Tissue Data

---

<b>Tissue Type:</b>	Serum
<b>Sex:</b>	Mixed
<b>Strain:</b>	Goat - Mixed

---

## Formulation

---

<b>Physical State:</b>	Liquid
<b>Concentration:</b>	102mg/ml by Refractometry
<b>Buffer:</b>	None
<b>Sterility:</b>	Non-sterile
<b>Preservative:</b>	None
<b>Stabilizer:</b>	None

---

## Shipping & Handling

---

<b>Shipping Condition:</b>	Dry Ice
----------------------------	---------

---

**Storage Condition:** Store container at -20° C prior to opening. Avoid cycles of freezing and thawing.

**Expiration:** Expiration date is one (1) year from date of receipt.

## Images



### Bottle

Normal Goat Serum can be used as a blocking agent to treat plastic surfaces, membrane or tissue after they have been sensitized with primary antibody or antigen. It provides an alternative to bovine serum albumin (BSA) and non-fat dry milk. It is effective in reducing nonspecific binding of proteins to reaction surfaces, thereby maximizing signal-to-noise ratio. This blocking agent is recommended for use in immunoassays where the primary antibody was produced in goat, as a source of non-specific serum protein or on tissue for immunohistochemical applications.

## References

- Nnah IC et al. TFEB-driven endocytosis coordinates MTORC1 signaling and autophagy. *Autophagy*. (2019)
- Kohno, D et al. Sweet Taste Receptor Serves to Activate Glucose- and Leptin-Responsive Neurons in the Hypothalamic Arcuate Nucleus and Participates in Glucose Responsiveness. *Frontiers in Neuroscience* (2016)
- Guo, L et al. Use of Human Induced Pluripotent Stem Cell-Derived Cardiomyocytes (hiPSC-CMs) to Monitor Compound Effects on Cardiac Myocyte Signaling Pathways. *Current Protocols in Chemical Biology* (2015)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.