



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet for KOA0235

**Mouse IL-12 (p70) ELISA Kit****Overview**

<b>Description:</b>	Mouse IL-12(p70) AccuSignal ELISA Kit - KOA0235
<b>Item No.:</b>	KOA0235
<b>Size:</b>	1 Kit
<b>Applications:</b>	ELISA
<b>Reactivity:</b>	Mouse
<b>Range:</b>	15.6 pg/ml - 1000 pg/ml

**Product Details**

<b>Background:</b>	Interleukin(IL)-12 is a 70-KDa cytokine comprised of two disulfide-linked proteins(p35 and p40) and is essential for the initiation of effective immune response. And the IL-12p70 is a heterodimer of p35 and p40 subunits; it is an important cytokine secreted by antigen-presenting cells in response to antigenic stimulation. Gene expression analysis of the IL-12 cytokine family subunits revealed that both strains induced high levels of p40(protein chain communal to IL-12 p70 and IL-23) as well as p19, a subunit of IL-23. Conversely only ACT- 18HS19 infection induced consistent transcription of IL-12 p35, a subunit of IL-12 p70. The standard product used in this kit is recombinant mouse IL-12 p70 comprising of a 193 amino acids p35 chain and a 313 amino acids p40 chain connected by a disulfide bond.
<b>Synonyms:</b>	CLMF, CLMF p35, Cytotoxic lymphocyte maturation factor 1, Cytotoxic lymphocyte maturation factor 1 p35, Cytotoxic lymphocyte maturation factor 35 kDa subunit, IL-12 subunit p35, IL-12A, IL12A, IL12A_HUMAN, IL35 subunit, Interleukin 12 alpha chain, Interleukin 12 p35, Interleukin 12 p70, Interleukin-12 subunit alpha, Natural killer cell stimulatory factor 1, Natural killer cell stimulatory factor 1 35 kD subunit, NF cell stimulatory factor chain 1, NFSK, NK cell stimulatory factor chain 1, NKSF1, P35
<b>Detection Kit Type:</b>	ELISA Kit
<b>Detection Range:</b>	15.6 pg/ml - 1000 pg/ml
<b>Sensitivity:</b>	<1.5 pg/ml

**Target Details**

<b>Gene Name:</b>	IL12A
-------------------	-------

<b>Reactivity:</b>	Mouse
<b>Immunogen:</b>	Expression system for standard: SF21; Immunogen sequence: (M23-S419)+(R23-A215)
<b>Purity/Specificity:</b>	Natural and recombinant mouse IL-12(P70). There is no detectable cross-reactivity with other relevant proteins.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - F8WI71</a></li><li>• <a href="#">NCBI - NP_001152896.1</a></li><li>• <a href="#">GenelD - 16159</a></li><li>• <a href="#">KOA0235 Protocol</a></li></ul>

## Application Details

<b>Tested Applications:</b>	ELISA
<b>Application Note:</b>	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml, 31.2pg/ml, 15.6pg/ml mouse IL-12(P70) standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of mouse cell culture supernates or serum to each empty well. It is recommended that each mouse IL-12(P70) standard solution and each sample be measured in duplicate.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	15.6 pg/ml-1000 pg/ml
<b>Other:</b>	DOE: 2-8° C -12/13/2024 -20° C - 06/13/2025

## Formulation


<b>Concentration:</b>	1 Kit
-----------------------	-------

## Shipping & Handling

<b>Shipping Condition:</b>	Wet Ice
<b>Storage Condition:</b>	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

**Expiration:** See kit insert for complete instructions.

## Images



### ELISA

Mouse IL-12(p70) AccuSignal ELISA Kit standard curve. Assay Range: 15.6pg/ml-1000pg/ml. Sensitivity: <1.5pg/ml.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.