

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





www.rockland.com tech@rockland.com +1 484.791.3823

#### Datasheet for KOA0262

## Mouse MIP-3 alpha - CCL20 ELISA Kit

#### **Overview**

Description:	Mouse MIP-3 alpha/CCL20 AccuSignal ELISA Kit - KOA0262
Item No.:	KOA0262
Size:	1 Kit
Applications:	ELISA
Reactivity:	Mouse
Range:	7.8 pg/ml - 500 pg/ml

#### **Product Details**

Background:	Macrophage Inflammatory Protein 3alpha(MIP3alpha), also called Chemokine, cc motif, ligand
	20/CCL20) The MID 2alpha/CCL20 gape was slened and sequenced revealing a four even three

20(CCL20). The MIP-3alpha/CCL20 gene was cloned and sequenced, revealing a four exon, three intron structure, and was localized by FISH analysis to 2q35-q36. MIP3alpha is predominantly expressed in lymph nodes, appendix, PBL, fetal liver, fetal lung and several cell lines. MIP3alpha/CCL20 and its receptor CCR6 are markedly up-regulated in psoriasis, and they may play a role in the recruitment of T cells to lesional psoriatic skin. And Alanine MIP-3alpha and Serine MIP-3alpha, the two forms of MIP3alpha, that differ by one amino acid at the predicted signal peptide cleavage site. Both of them were chemically synthesized and tested for biological activity. And both flu antigen plus IL-2-activated CD4(+) and CD8(+) T lymphoblasts and cord

blood-derived dendritic cells responded to Ser and Ala MIP-3alpha.

**Synonyms:** Beta chemokine exodus 1, Beta-chemokine exodus-1, C C motif chemokine 20, CC chemokine

LARC, Ccl20, CCL20(2-70), CCL20\_HUMAN, Chemokine (C C motif) ligand 20, CKb4, Exodus 1, LARC, Liver and activation regulated chemokine, Liver and activation-regulated chemokine, Macrophage inflammatory protein 3 alpha, MIP 3 alpha, MIP 3a, MIP-3-alpha, MIP3 alpha, MIP3A, MIP3Alpha, SCYA20, Small inducible cytokine A20, Small inducible cytokine subfamily A

(Cys Cys) member 20, Small-inducible cytokine A20, ST38

**Detection Kit Type:** ELISA Kit

**Detection Range:** 7.8 pg/ml - 500 pg/ml

Sensitivity: <1 pg/ml

### **Target Details**

Gene Name: CCL20

www.rockland.com Page 1 of 3





www.rockland.com tech@rockland.com +1 484.791.3823

Reactivity:	Mouse
Immunogen:	Expression system for standard: E.coli; Immunogen sequence: A28-M97
Purity/Specificity:	Natural and recombinant mouse CCL20. There is no detectable cross-reactivity with other relevant proteins.
Relevant Links:	• UniProtKB - 089093
	• NCBI - NP_001153210.1
	• GeneID - 20297
	KOA0262 Protocol

## **Application Details**

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml, 31.3pg/ml, 15.6pg/ml, 7.8pg/ml mouse CCL20 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. We recommend that each mouse CCL20 standard solution and each sample is measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	DOE: 04/18/20 at 4°C 10/18/20 at -20°C

### **Formulation**

Concentration:	1 Kit
Anticoagulant:	Heparin Sodium

## **Shipping & Handling**

<b>Shipping Condition:</b>	Wet Ice
Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

www.rockland.com Page 2 of 3






www.rockland.com tech@rockland.com +1 484.791.3823

**Expiration:** See kit insert for complete instructions.

### **Images**



#### **ELISA**

Mouse MIP-3 alpha/CCL20 AccuSignal ELISA kit standard curve. Assay Range: 7.8pg/ml-500pg/ml. Sensitivity: <1pg/ml.

#### **Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com Page 3 of 3