



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



HIP2 (h): 293T Lysate: sc-116355

BACKGROUND

HIP1 (Huntingtin interacting protein 1), a membrane-associated protein and HIP2 bind specifically to the N-terminus of human Huntingtin. HIP1 and HIP2 are ubiquitously expressed in different brain regions at low levels and exhibit nearly identical subcellular fractionation as Huntington. The Huntington-HIP1 interaction is inversely correlated to the polyglutamine length in Huntington, suggesting that loss of normal Huntington-HIP1 interaction may compromise the membrane-cytoskeletal integrity in the brain. Conversely, the Huntington-HIP2 interaction is not affected by the polyglutamine length in the Huntington protein. However, both HIP1 and HIP2 play an important role in the pathogenesis of Huntington disease (HD).

REFERENCES

1. Sun, H., et al. 1992. Effects of McAbs HIP2, APT4 and HI117 on the human platelet cytoskeleton. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao* 14: 1-5.
2. Kalchman, M.A., et al. 1996. Huntington is ubiquitinated and interacts with a specific ubiquitin-conjugating enzyme. *J. Biol. Chem.* 271: 19385-19394.
3. Tanno, Y., et al. 1999. Localization of Huntington-interacting protein-2 (HIP2) mRNA in the developing mouse brain. *J. Chem. Neuroanat.* 17: 99-107.
4. Wang, Y., et al. 2000. YAC/BAC-based physical and transcript mapping around the gracile axonal dystrophy (GAD) locus identifies UCH-L1, Pmx2b, Atp3a2 and HIP2 genes. *Genomics* 66: 333-336.
5. Lee, S.J., et al. 2001. E3 ligase activity of RING finger proteins that interact with HIP2, a human ubiquitin-conjugating enzyme. *FEBS Lett.* 503: 61-64.
6. Song, S., et al. 2003. Essential role of E2-25K/HIP2 in mediating Amyloid- β neurotoxicity. *Mol. Cell* 12: 553-563.
7. Wesierska-Gadek, J., et al. 2007. Roscovitine-activated HIP2 kinase induces phosphorylation of wt p53 at Ser 46 in human MCF7 breast cancer cells. *J. Cell. Biochem.* 100: 865-874.
8. Metzler, M., et al. 2007. NMDA receptor function and NMDA receptor-dependent phosphorylation of Huntington is altered by the endocytic protein HIP1. *J. Neurosci.* 27: 2298-2308.
9. Ybe, J.A., et al. 2007. Crystal structure at 2.8 Å of the DLLRKN-containing coiled-coil domain of Huntington-interacting protein 1 (HIP1) reveals a surface suitable for clathrin light chain binding. *J. Mol. Biol.* 367: 8-15.

CHROMOSOMAL LOCATION

Genetic locus: UBE2K (human) mapping to 4p14.

PRODUCT

HIP2 (h): 293T Lysate represents a lysate of human HIP2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

HIP2 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive HIP2 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


HIP2 (H-6): sc-390339 is recommended as a positive control antibody for Western Blot analysis of enhanced human HIP2 expression in HIP2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



HIP2 (H-6): sc-390339. Western blot analysis of HIP2 expression in non-transfected: sc-117752 (**A**) and human HIP2 transfected: sc-116355 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.