



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# CENP-B (h): 293T Lysate: sc-116535

## BACKGROUND

A replicated chromosome includes two kinetochores that control chromosome segregation during mitosis. Both centromere proteins CENP-B and CENP-H are contained in the centromeric heterochromatin between kinetochores, and are involved in maintaining sister chromatid cohesion. The highly dispersed CENP-B promotes and maintains the joining of DNA satellites in the centromere. CENP-B targets centromeric  $\alpha$ -DNA and protects it from digestion by nucleases as well as preventing DNase or restriction enzyme digestion from affecting the morphology of centromeres. CENP-H contains a coiled-coil structure and a nuclear localization signal. CENP-H is specifically and constitutively localized to kinetochores and plays a role in the organization and function of kinetochores throughout the cell cycle.

## REFERENCES

1. Cooke, C.A., et al. 1990. CENP-B: a major human centromere protein located beneath the kinetochore. *J. Cell Biol.* 110: 1475-1488.
2. Rieder, C.L., et al. 1998. The vertebrate cell kinetochore and its roles during mitosis. *Trends Cell Biol.* 8: 310-318.
3. Barbosa-Cisneros, O., et al. 1998. Localization of the centromere protein CENP-B using scleroderma sera and evidence for a role in centromere survival. *Rev. Rhum. Engl. Ed.* 65: 15-20.
4. Sugata, N., et al. 1999. Characterization of a novel kinetochore protein, CENP-H. *J. Biol. Chem.* 274: 27343-27346.
5. Choo, K.H. 2000. Centromerization. *Trends Cell Biol.* 10: 182-188.
6. Ohzeki, J., et al. 2002. CENP-B box is required for *de novo* centromere chromatin assembly on human alphoid DNA. *J. Cell Biol.* 159: 765-775.
7. Suzuki, N., et al. 2004. CENP-B interacts with CENP-C domains containing Mif2 regions responsible for centromere localization. *J. Biol. Chem.* 279: 5934-5946.
8. Lomonte, P. et al. 2007. Centromeric protein CENP-B proteasomal degradation induced by the viral protein ICP0. *FEBS Lett.* 581: 658-662.
9. Dolnik, A.V. et al. 2007. Dynamics of satellite binding protein CENP-B and telomere binding protein TRF2/MTBP in the nuclei of mouse spermatogenic line. *Cell Biol Int.* 31: 316-329

## CHROMOSOMAL LOCATION

Genetic locus: CENPB (human) mapping to 20p13.

## PRODUCT

CENP-B (h): 293T Lysate represents a lysate of human CENP-B transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

CENP-B (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive CENP-B antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


CENP-B (C-10): sc-376392 is recommended as a positive control antibody for Western Blot analysis of enhanced human CENP-B expression in CENP-B transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



CENP-B (C-10): sc-376392. Western blot analysis of CENP-B expression in non-transfected: sc-117752 (**A**) and human CENP-B transfected: sc-116535 (**B**) 293T whole cell lysates.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.