



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Mel-CAM (h): 293T Lysate: sc-116616

## BACKGROUND

The tumorigenic and metastatic phenotype of melanoma cells correlates well with an increased expression of cell-cell and cell-matrix adhesion receptors. The human Mel-CAM gene maps to chromosome 11q23 and encodes a transmembrane glycoprotein, also designated MCAM, MUC18 or CD146, that belongs to the immunoglobulin superfamily and functions as a  $Ca^{2+}$ -independent cell adhesion molecule. The deduced human sequence of 603 amino acids consists of a signal peptide, five immunoglobulin-like domains, a transmembrane region and a short cytoplasmic tail. Mel-CAM expression is restricted to advanced primary and metastatic melanomas and to cell lines of the neuroectodermal lineage, but not normal melanocytes. Mel-CAM is found on 80% of advanced primary human melanomas and correlates well with development of metastatic disease. Mel-CAM activation initiates an outside-in signaling pathway that involves the protein tyrosine kinases Fyn, FAK and paxillin. Mel-CAM influences the dynamics of Actin cytoskeleton rearrangement and is essential for the maintenance of thymic architecture and function.

## REFERENCES

1. Lehmann, J.M., et al. 1989. MUC18, a marker of tumor progression in human melanoma, shows sequence similarity to the neural cell adhesion molecules of the immunoglobulin superfamily. *Proc. Natl. Acad. Sci. USA* 86: 9891-9895.
2. Kuzu, I., et al. 1993. Expression of adhesion molecules on the endothelium of normal tissue vessels and vascular tumors. *Lab. Invest.* 69: 322-328.
3. Sers, C., et al. 1993. Genomic organization of the melanoma-associated glycoprotein MUC18: implications for the evolution of the immunoglobulin domains. *Proc. Natl. Acad. Sci. USA* 90: 8514-8518.
4. Vainio, O., et al. 1996. HEMCAM, an adhesion molecules expressed by c-Kit<sup>+</sup> hemopoietic progenitors. *J. Cell Biol.* 135: 1655-1668.
5. Shih, I.M. 1999. The role of CD146 (Mel-CAM) in biology and pathology. *J. Pathol.* 189: 4-11.
6. Seftalioglu, A. and Karakoc, L. 2000. Expression of CD146 adhesion molecules (MUC18 or MCAM) in the thymic microenvironment. *Acta Histochem.* 102: 69-83.

## CHROMOSOMAL LOCATION

Genetic locus: MCAM (human) mapping to 11q23.3.

## PRODUCT

Mel-CAM (h): 293T Lysate represents a lysate of human Mel-CAM transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS

Mel-CAM (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Mel-CAM antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Mel-CAM (2Q401): sc-71565 is recommended as a positive control antibody for Western Blot analysis of enhanced human Mel-CAM expression in Mel-CAM transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Mel-CAM (2Q401): sc-71565. Western blot analysis of Mel-CAM expression in non-transfected: sc-117752 (A) and human Mel-CAM transfected: sc-116616 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.