



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Synaptotagmin I (h): 293T Lysate: sc-116669

BACKGROUND

Synaptotagmins are a large gene family of synaptic vesicle type III integral membrane proteins that function as regulators of both exocytosis and endocytosis, and are involved in neurotransmitter secretion from small secretory vesicles. Calcium binds to Synaptotagmin I, which triggers neurotransmitter release at the synapse. Synaptotagmin II is phosphorylated by WNK1 in a process that regulates calcium-dependent interactions. Synaptotagmin III is involved in calcium-dependent exocytosis of secretory vesicles in endocrine cells and neurons. Synaptotagmin IV is expressed in neuronal tissues, and has the highest mRNA levels in the hippocampus. The proximity of the Synaptotagmin IV gene to markers of several psychiatric disorders suggest an involvement of synaptotagmin IV in human disease. Synaptotagmin V is a dense-core vesicle-specific protein that regulates a specific type of calcium-regulated secretion. Synaptotagmin VI interacts with adaptor protein-2 in a calcium-independent manner. Synaptotagmin VII is widely expressed in non-neuronal tissues.

REFERENCES

- Hilbush, B.S. and Morgan, J.I. 1994. A third synaptotagmin gene, Syt3, in the mouse. *Proc. Natl. Acad. Sci. USA* 91: 8195-8199.
- Li, C., Ullrich, B., Zhang, J.Z., Anderson, R.G., Brose, N. and Sudhof, T.C. 1995. Ca²⁺-dependent and -independent activities of neural and non-neuronal Synaptotagmins. *Nature* 375: 594-599.
- Kishore, B.K., Wade, J.B., Schorr, K., Inoue, T., Mandon, B. and Knepper, M.A. 1998. Expression of Synaptotagmin VIII in rat kidney. 1998. Expression of Synaptotagmin VIII in rat kidney. *Am. J. Physiol.* 275: 131-142.
- Xi, D., Chin, H. and Gainer, H. 1999. Analysis of Synaptotagmin I-IV messenger RNA expression and developmental regulation in the rat hypothalamus and pituitary. *Neuroscience* 88: 425-435.
- Ferguson, G.D., Chen, X.N., Korenberg, J.R. and Herschman, H.R. 2000. The human Synaptotagmin IV gene defines an evolutionary break point between syntenic mouse and human chromosome regions but retains ligand inducibility and tissue specificity. *J. Biol. Chem.* 275: 36920-3696.
- LocusLink Report (LocusID: 6859). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: SYT1 (human) mapping to 12q21.2.

PRODUCT

Synaptotagmin I (h): 293T Lysate represents a lysate of human Synaptotagmin I transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Synaptotagmin I (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Synaptotagmin I antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


Synaptotagmin I/II (H-9): sc-393392 is recommended as a positive control antibody for Western Blot analysis of enhanced human Synaptotagmin I expression in Synaptotagmin I transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG_x BP-HRP: sc-516102 or m-IgG_x BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Synaptotagmin I/II (H-9): sc-393392. Western blot analysis of Synaptotagmin I expression in non-transfected: sc-117752 (**A**) and human Synaptotagmin I transfected: sc-116669 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.